

R 9  
45 92

DEC. 9  
HIC  
239

MTP-P&VE-F-62-12  
December 12, 1962

NASA TMX 50,106

N 63 17342

**GEORGE C. MARSHALL**

**SPACE  
FLIGHT  
CENTER**

**HUNTSVILLE, ALABAMA**

**On a Consistent System of  
Astrodynamical  
Constants**

By

*Helmut G. L. Krause*



**FOR INTERNAL USE ONLY**

**AERONAUTICS AND SPACE ADMINISTRATION**

RQT  
4597

Copy A

Welle

GEORGE C. MARSHALL SPACE FLIGHT CENTER

MTP-P&VE-F-62-12

ON A CONSISTENT SYSTEM OF ASTRODYNAMIC CONSTANTS

*By*

*Helmut G. L. Krause*


**ABSTRACT**

An internally consistent system of astrodynamical constants is derived based upon theoretical coupling relationships and the most recent available experimental data. A previously existing discrepancy in the value of the gravitational parameter of the earth (as derived by different methods) has been eliminated. Likewise, several inconsistencies in the previously available system of lunar constants have been removed.

A new method of determining the ratio of the masses of the Earth and Moon has been derived and the results are in agreement with other determinations.

An error study of each constant is presented; both relative and absolute probable errors are listed.

*NASA-TM-X-50,106*



GEORGE C. MARSHALL SPACE FLIGHT CENTER

MTP-P&VE-F-62-12

ON A CONSISTENT SYSTEM OF ASTRODYNAMIC CONSTANTS

*By*


*Helmut G. L. Krause*

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

GEORGE C. MARSHALL SPACE FLIGHT CENTER

PROPULSION AND VEHICLE ENGINEERING DIVISION

HUNTSVILLE, ALABAMA



## TABLE OF CONTENTS

<u>Section</u>		<u>Page</u>
Summary (A List of Astrodynamical Constants) -----	1	
1. The Astronomical Unit and the Solar Parallax -----	11	
2. Definition of Time Units, Mean Orbital Motions, and Rotational Angular Velocities of Earth and Moon -----	12	
3. The Lunar Distance and the Lunar Parallax -----	17	
4. Mass Ratios of the Sun and the Earth-Moon System -----	18	
5. The Constant of Lunar Inequality and the Parallactic Inequality in Moon's Ecliptic Longitude -----	19	
6. Another Method for the Determination of the Ratio of the Masses of Earth and Moon -----	20	
7. Geodynamic (Terrestrial) Relations -----	22	
8. Experimental Values of the Gravitational Acceleration of the Earth -----	26	
9. The Dynamic Oblateness and the Constants of Precession and Nutation --	28	
10. Density Distribution Within the Moon -----	30	
11. Constants of the Physical Libration of the Moon -----	31	
12. Relations and Numerical Values for the Dimensionless Moment of Inertia Parameters -----	32	
13. The Final Determination of the Dimension, Masses and Moments of Inertia for the Earth and the Moon -----	34	
14. The Earth Ellipsoid -----	36	
References -----	45	
Approval -----	49	
Distribution -----	50	



GEORGE C. MARSHALL SPACE FLIGHT CENTER

MTP-P&VE-F-62-12

ON A CONSISTENT SYSTEM OF ASTRODYNAMIC CONSTANTS

*By*

*Helmut G. L. Krause*

**SUMMARY**

An internally consistent system of astrodynamical constants is derived based upon theoretical coupling relationships and the most recent available experimental data. A previously existing discrepancy in the value of the gravitational parameter of the earth (as derived by different methods) has been eliminated. Likewise, several inconsistencies in the previously available system of lunar constants have been removed.

A new method of determining the ratio of the masses of the Earth and Moon has been derived and the results are in agreement with other determinations.

An error study of each constant is presented; both relative and absolute probable errors are listed.

The results of this study can be summarized in the following list of astrodynamical constants:

## 1. GENERAL CONSTANTS

1. Newton's gravitation constant:

$$G = 6.670 (1 \pm 0.0007) \times 10^{-8} = (6.670 \pm 0.005) \times 10^{-8} \text{ cm}^3 \text{ g}^{-1} \text{ s}^{-2} (\text{dyn cm}^2 \text{ g}^{-2})$$

2. Velocity of light:

$$c = 299\,792.5 (1 \pm 3.3 \times 10^{-7}) = 299\,792.5 \pm 0.1 \text{ km/s}$$

3. Solar parallax (Sun's equatorial horizontal parallax):

$$\pi_{\odot} = 8.''794\,14 (1 \pm 5.8 \times 10^{-6}) = 8.''794\,14 \pm 0.''000\,05$$

4. Astronomical unit (mean Earth-Sun distance =  $R_{\odot} / \pi_{\odot} \sin 1''$ ):

$$a. \text{ u.} = 149\,598\,700 (1 \pm 2.7 \times 10^{-6}) = 149\,598\,700 \pm 400 \text{ km}$$

5. Light year (distance which light travels in a year =  $P_{\oplus} c$ ):

$$1. \text{ y.} = 31\,556\,925.9747 \text{ c} = (9.460\,530 \pm 0.000\,003) \times 10^{12} \text{ km} = 63\,239.39 \pm 0.15 \text{ a. u.}$$

6. Parsec (distance in which 1 a. u. appears as  $1'' = 1 \text{ a. u.} / \sin 1''$ ):

$$pc = 206\,264.806\,247 \text{ a. u.} = (3.085\,695 \pm 0.000\,008) \times 10^{13} \text{ km} = 3.261\,651 \pm 0.000\,008 \text{ l. y.}$$

7. Light time for 1 a. u. :

$$\tau = a.u./c = 499.008 (1 \pm 3.6 \times 10^{-5}) = 499.008 \pm 0.018 \text{ s}$$

8. Constant of aberration:

$$K = 20.''4956 (1 \pm 3.5 \times 10^{-5}) = 20.''4956 \pm 0.''0007$$

9. Obliquity of ecliptic

$$\epsilon = 23^\circ 27' 8.''26 - 46.''844 T - 0.''0060 T^2 + 0.''00183 T^3$$

$$\cos \epsilon = 0.917\,3917 ; \sin \epsilon = 0.397\,9855 (1900.0)$$

10. Newcomb's constant of precession (per tropical century):

$$P = \frac{p_0}{\cos \epsilon} = 5493.''62 - 0.''003\,64 T = (N + 2.''96)*$$

11. Luni-solar precession in longitude:

$$p_0 = p_{\odot} + p_{\oplus} = 5039.''804 + 0.''4930 T - 0.''000\,04 T^2 = (N + 2.''72)$$

\* N refers to Newcomb's precessional data

12. Geodetic precession in longitude:

$$p_g = \frac{3}{2} (\nu_{\oplus}/c)^2 n_{\oplus} = 1.^{\prime\prime}9188 \pm 0.^{\prime\prime}0002 = (N + 1.^{\prime\prime}92)$$

13. Observed luni-solar precession in longitude:

$$p_1 = p_0 - p_g = 5037.^{\prime\prime}885 + 0.^{\prime\prime}4930 T - 0.^{\prime\prime}00004 T^2 = (N + 0.^{\prime\prime}80)$$

14. Planetary precession in right ascension:

$$\lambda = -m + n \cot \epsilon = 12.^{\prime\prime}473 - 1.^{\prime\prime}8870 T - 0.^{\prime\prime}00014 T^2 = (N + 0.^{\prime\prime}00)$$

15. General precession in longitude:

$$p = p_1 - \lambda \cos \epsilon = m \cos \epsilon + n \sin \epsilon = 5026.^{\prime\prime}441 + 2.^{\prime\prime}2229 T + 0.^{\prime\prime}00026 T^2 = (N + 0.^{\prime\prime}80)$$

16. General precession in right ascension:

$$\begin{aligned} m = p_1 \cos \epsilon - \lambda &= 4609.^{\prime\prime}236 + 2.^{\prime\prime}7945 T + 0.^{\prime\prime}00012 T^2 = (N + 0.^{\prime\prime}73) \\ &= 307.^{\circ}2824 + 0.^{\circ}18630 + 0.^{\circ}000008 T^2 = (N + 0.^{\circ}0487) \end{aligned}$$

17. General precession in declination:

$$\begin{aligned} n = p_1 \sin \epsilon &= 2005.^{\prime\prime}005 - 0.^{\prime\prime}8533 T - 0.^{\prime\prime}00037 T^2 = (N + 0.^{\prime\prime}32) \\ &= 133.^{\circ}6670 - 0.^{\circ}05689 T - 0.^{\circ}000025 T^2 = (N + 0.^{\circ}0213) \end{aligned}$$

18. Mean sidereal time rate (for 1950.0):

$$\begin{aligned} \dot{\Theta}_m &= 7.292\,115\,854\,79 \times 10^{-5} \text{ rad/s} = 1.002\,737\,909\,294 \text{ d}_*/\text{d} (\text{s}_*/\text{s}) \\ &= 15.041\,068\,639\,41 ''/\text{s} (\text{°}/\text{h}) = 360.985\,647\,346\,0 \text{ °}/\text{d} \end{aligned}$$

## 2. EARTH CONSTANTS

19. Semi-major axis of the Earth's orbit:

$$a_{\oplus} = 149\,598\,700 (1 \pm 2.7 \times 10^{-6}) = 149\,598\,700 \pm 400 \text{ km}$$

20. Siderial mean orbital motion (for 1950.0):

$$\begin{aligned} n_{\oplus} &= 0.985\,609\,108\,0 \text{ °}/\text{d} = 0.041\,067\,046\,15 ''/\text{s} (\text{°}/\text{h}) \\ &= 1.990\,986\,581\,7 \times 10^{-7} \text{ rad/s} \end{aligned}$$

21. Mean orbital velocity:

$$v_{\oplus} = a_{\oplus} n_{\oplus} = 29784.90 (1 \pm 2.7 \times 10^{-6}) = 29784.90 \pm 0.08 \text{ m/s}$$

22. Mass ratio of the Sun to the Earth-Moon system:

$$\nu = \frac{M_{\odot}}{M_{\oplus} + M_{\odot}} = 328898.6 (1 \pm 1.6 \times 10^{-5}) = 328898.6 \pm 5.2$$

23. Mass ratio of the Sun to the Earth:

$$\nu (1 + \kappa) = \frac{M_{\odot}}{M_{\oplus}} = 332947.6 (1 \pm 2.0 \times 10^{-5}) = 332947.6 \pm 6.7$$

24. Gravitational parameter of the Earth:

$$\mu_{\oplus} = GM_{\oplus} = 398606.4 (1 \pm 1.23 \times 10^{-5}) = 398606.4 \pm 4.9 \text{ km}^3/\text{s}^2$$

25. Mass:

$$M_{\oplus} = 5.9761 \times 10^{27} (1 \pm 7.2 \times 10^{-4}) = (5.9761 \pm 0.0043) \times 10^{27} \text{ g}$$

26. Equatorial radius:

$$R_{\oplus} = 6378170 (1 \pm 3.14 \times 10^{-6}) = 6378170 \pm 20 \text{ m}$$

27. Polar radius:

$$R_p = 6356788 (1 \pm 3.70 \times 10^{-6}) = 6356788 \pm 24 \text{ m}$$

28. Flattening (oblateness, ellipticity):

$$f = \frac{R_{\oplus} - R_p}{R_{\oplus}} = 0.00335233 (1 \pm 1.7 \times 10^{-4}) = 0.00335233 \pm 0.00000056 = 1 : (298.30 \pm 0.05)$$

$$1 - f = (1 - e^2)^{1/2} = (1 + \epsilon^2)^{-1/2} \approx R_p / R_{\oplus} = 0.99664767 (1 \pm 5.6 \times 10^{-7})$$

29. First eccentricity of the meridian ellipse:

$$e = 0.08181333 + 0.00000680$$

$$e^2 = f(2 - f) = 0.00669342 + 0.00000111$$

30. Second eccentricity of the meridian ellipse:

$$\epsilon = 0.08208852 \pm 0.00000687$$

$$\epsilon^2 = \frac{e^2}{1 - e^2} = 0.00673853 \pm 0.00000113$$

31. Mean radius:

$$\bar{R} = (2R_{\oplus} + R_p) / 3 = 6371043 (1 \pm 3.3 \times 10^{-6}) = 6371043 \pm 21 \text{ m}$$

32. Radius for geodetic latitude  $\varphi = \sin^{-1} \sqrt{1/3} = 35^\circ 15' 51.''8$

$$R_1 = 6371\,083 (1 \pm 3.3 \times 10^{-6}) = 6371\,083 \pm 21 \text{ m}$$

33. Radius for geocentric latitude  $\phi = \sin^{-1} \sqrt{1/3} = 35^\circ 15' 51.''8$

$$R_2 = 6371\,019 (1 \pm 3.3 \times 10^{-6}) = 6371\,019 \pm 21 \text{ m}$$

34. Radius for sphere of same surface area:

$$R_s = 6371\,041 (1 \pm 3.3 \times 10^{-6}) = 6371\,041 \pm 21 \text{ m}$$

35. Radius for sphere of same volume:

$$R_v = 6371\,035 (1 \pm 3.3 \times 10^{-6}) = 6371\,035 \pm 21 \text{ m}$$

36. Surface area:

$$S_\oplus = 5.100\,711 \times 10^{14} (1 \pm 6.6 \times 10^{-6}) = (5.100\,711 \pm 0.000\,034) \times 10^{14} \text{ m}^2$$

37. Volume:

$$V_\oplus = 1.083\,225 \times 10^{21} (1 \pm 1.0 \times 10^{-5}) = (1.083\,225 \pm 0.000\,011) \times 10^{21} \text{ m}^3$$

38. Mean density:

$$\bar{\rho}_\oplus = 5.5170 (1 \pm 7.3 \times 10^{-4}) = 5.5170 \pm 0.0040 \text{ g/cm}^3$$

39. Angular velocity of the Earth's rotation:

$$\begin{aligned} \Omega_m &= 7.292\,115\,146\,46 \times 10^{-5} \text{ rad/s} = 1.002\,737\,811\,891 \text{ rot/d} \\ &= 15.041\,067\,178\,37 \text{ ''/s } (\text{°}/\text{h}) = 360.985\,612\,280.8 \text{ °/d} \end{aligned}$$

40. Rotational velocity at the equator:

$$\Omega_m R_e = 465.1035 (1 \pm 3.2 \times 10^{-6}) = 465.1035 \pm 0.0015 \text{ m/s}$$

41. Centrifugal acceleration at the equator:

$$\Omega_m^2 R_e = 0.033\,915\,88 (1 \pm 3.2 \times 10^{-6}) = 0.033\,915\,88 \pm 0.000\,000\,11 \text{ m/s}^2$$

42. Centrifugal acceleration factor:

$$\tilde{\omega}_\oplus = \frac{\Omega_m^2 R_e^3}{G M_\oplus} = 3461.369 \times 10^{-6} (1 \pm 2.2 \times 10^{-5}) = (3461.369 \pm 0.076) \times 10^{-6}$$

43. Oblateness coefficients of the Earth's potential:

$$J = 1623.48 \times 10^{-6} (1 \pm 1.8 \times 10^{-4}) = (1623.48 \pm 0.29) \times 10^{-6}$$

$$K = \frac{6}{7} \quad D = 8.85 \times 10^{-6}$$

$$J_2 = \frac{2}{3} \quad J = 1082.32 \times 10^{-6} (1 \pm 1.8 \times 10^{-4}) = (1082.32 \pm 0.19) \times 10^{-6}$$

$$J_4 = -\frac{4}{15} \quad K = -\frac{8}{35} \quad D = -2.36 \times 10^{-6}$$

44. Coefficients of the Earth's gravity formula:

$$\beta = 5302.92 \times 10^{-6}; \quad \gamma = -5.85 \times 10^{-6}$$

45. Mass of the Earth's atmosphere:

$$M_{\text{atm}} = \frac{p_0}{g} \cdot S_{\oplus} = (10332.275 \text{ kg/m}^2) \cdot S_{\oplus} \\ = 5.270195 \times 10^{18} (1 \pm 6.6 \times 10^{-6}) = (5.270195 \pm 0.000035) \times 10^{18} \text{ kg}$$

46. Relative mass of the Earth's atmosphere:

$$A = M_{\text{atm}} / M_{\oplus} = 0.88188 \times 10^{-6} (1 \pm 7.3 \times 10^{-4}) = (0.88188 \pm 0.00064) \times 10^{-6}$$

47. Gravity acceleration correction factor:

$$\chi = g_e / (GM_{\oplus} / R_e^2) = 0.99816566 (1 \pm 4.0 \times 10^{-7}) = 0.99816566 \pm 0.00000040$$

$$1 - \chi = A + \tilde{\omega} - J - \frac{1}{2} K = 1834.34 \times 10^{-6} (1 \pm 2.2 \times 10^{-4}) = (1834.34 \pm 0.40) \times 10^{-6}$$

48. Gravity acceleration at the Earth's equator:

$$g_e = 9.780362 (1 \pm 3.3 \times 10^{-6}) = 9.780362 \pm 0.000032 \text{ m/s}^2$$

49. Dynamic oblateness:

$$H = \frac{C - A}{C} = 3272.09 \times 10^{-6} (1 \pm 1.6 \times 10^{-4}) = (3272.09 \pm 0.54) \times 10^{-6} = 1/305.615 \pm 0.05$$

50. Moment of inertia parameter:

$$q = \frac{3}{2} \cdot \frac{C}{M_{\oplus} R_e^2} = \frac{J}{H} = 0.49616 (1 \pm 3.4 \times 10^{-4}) = 0.49616 \pm 0.00017$$

51. Dimensionless moments of inertia:

$$\frac{A}{M_{\oplus} R_e^2} = J_2 \left( \frac{1}{H} - 1 \right) = 0.32969 \pm 0.00011$$

$$\frac{C}{M_{\oplus} R_e^2} = J_2 / H = 0.33077 \pm 0.00011$$

52. Unit for the Earth's moments of inertia:

$$M_{\oplus} R_e^2 = 2.4311_4 \times 10^{38} (1 \pm 7.3 \times 10^{-4}) = (2.4311_4 \pm 0.0018) \times 10^{38} \text{ kg m}^2$$

53. Earth's moments of inertia:

$$A = 0.80152 \times 10^{38} (1 \pm 1.07 \times 10^{-3}) = (0.80152 \pm 0.00086) \times 10^{38} \text{ kg m}^2$$

$$C = 0.80415 \times 10^{38} (1 \pm 1.07 \times 10^{-3}) = (0.80415 \pm 0.00086) \times 10^{38} \text{ kg m}^2$$

$$C - A = 2.6313 \times 10^{35} (1 \pm 9.1 \times 10^{-4}) = (2.6313 \pm 0.0024) \times 10^{35} \text{ kg m}^2$$

54. Angular momentum:

$$C \Omega = 5.8640 \times 10^{33} (1 \pm 1.07 \times 10^{-3}) = (5.8640 \pm 0.0063) \times 10^{33} \text{ kg m}^2/\text{s}$$

55. Rotational energy:

$$\frac{1}{2} C \Omega^2 = 2.1380 \times 10^{29} (1 \pm 1.07 \times 10^{-3}) = 2.1380 \pm 0.0023 \times 10^{29} \text{ kg m}^2/\text{s}^2 (\text{joule})$$

56. Circular velocity at the Earth's equator:

$$v_{\text{cir}} = \sqrt{\mu_{\oplus}/R_e} = 7905.404 (1 \pm 7.7 \times 10^{-6}) = 7905.404 \pm 0.061 \text{ m/s}$$

### 3. LUNAR CONSTANTS

57. Mean observed distance from the Earth:

$$\bar{r}_e = 384402.0 (1 \pm 2.6 \times 10^{-6}) = 384402.0 \pm 1.0 \text{ km}$$

58. Relative mean lunar distance:

$$\bar{r}_e/R_e = 1/\sin \pi_e = 60.26838 (1 \pm 5.8 \times 10^{-6}) = 60.26838 \pm 0.00035$$

59. Constant part of the sine of the perturbed lunar parallax:

$$\pi'_e = \frac{\sin \pi_e}{\sin 1} = 3422.^{\prime\prime}438 (1 \pm 5.8 \times 10^{-6}) = 3422.^{\prime\prime}438 \pm 0.^{\prime\prime}020$$

60. Mean perturbed equatorial horizontal parallax:

$$\pi_e = 3422.^{\prime\prime}595 (1 \pm 5.8 \times 10^{-6}) = 3422.^{\prime\prime}595 \pm 0.^{\prime\prime}020$$

61. Semi-major axis of the Moon's orbit:

$$a_e = 1.000907681 \quad \bar{r}_e = 384750.9 \quad (1 \pm 2.6 \times 10^{-6}) = 384750.9 \pm 1.0 \text{ km}$$

62. Siderial mean orbital motion (for 1950.0):

$$\pi_e = 13.1763582598 ^{\circ}/d = 0.549014912685 ^{\prime\prime}/s \quad (^{\circ}/h)$$

$$= 2.66169940799 \times 10^{-6} \text{ rad/s}$$

63. Mean orbital velocity:

$$v_e = a_e n_e = 1024.091 (1 \pm 2.9 \times 10^{-6}) = 1024.091 \pm 0.003 \text{ m/s}$$

64. Lunar inequality in the Moon's ecliptic longitude:

$$L = 6.''4439 (1 \pm 3.0 \times 10^{-4}) = 6.''4439 \pm 0.''0019$$

65. Parallactic inequality in the Moon's ecliptic longitude:

$$P_* = 124.''986 (1 \pm 3.3 \times 10^{-5}) = 124.''986 \pm 0.''004$$

66. Mass ratio of the Earth to the Moon:

$$1/\kappa = M_\oplus / M_e = 81.250 (1 \pm 3.0 \times 10^{-4}) = 81.250 \pm 0.024$$

67. Mass ratio of the Earth-Moon system to the Earth:

$$1 + \kappa = (M_\oplus + M_e) / M_\oplus = 1.012\,307\,7 (1 \pm 3.7 \times 10^{-6}) = 1.012\,307\,7 \pm 0.000\,0037$$

68. Gravitational parameter of the Moon:

$$\mu_e = GM_e = 4905.92 (1 \pm 3.1 \times 10^{-4}) = 4905.92 \pm 1.52 \text{ km}^3/\text{s}^2$$

69. Mass:

$$M_e = 7.3552 \times 10^{25} (1 \pm 1.02 \times 10^{-3}) = (7.3552 \pm 0.0075) \times 10^{25} \text{ g}$$

70. Moon's semi-diameter at mean distance:

$$s_e = 932.''72 (1 \pm 1.0 \times 10^{-4}) = 932.''72 \pm 0.''09$$

71. Relative radius of the visible disk of the Moon:

$$k = R_e / R_\oplus = 0.2725\,289 (1 \pm 1.0 \times 10^{-4}) = 0.2725\,289 \pm 0.0000\,273$$

72. Radius of the visible disk of the Moon:

$$R_e = \frac{b+c}{2} = 1738\,236 (1 \pm 1.0 \times 10^{-4}) = 1738\,236 \pm 174 \text{ m}$$

73. Longest semi-axis directed to the Earth:

$$a = R_e / 0.9995\,918 = 1738\,946 (1 \pm 1.07 \times 10^{-4}) = 1738\,946 \pm 186 \text{ m}$$

74. Medium semi-axis in orbital direction:

$$b = 0.9998\,116 \quad a = 1738\,618 (1 \pm 1.20 \times 10^{-4}) = 1738\,618 \pm 209 \text{ m}$$

75. Shortest semi-axis (rotational or polar radius):

$$c = 0.9993\,720 \quad a = 1737\,854 (1 \pm 1.08 \times 10^{-4}) = 1737\,854 \pm 188 \text{ m}$$

76. Volume:

$$V = \frac{4}{3} \pi a b c = 2.20086 \times 10^{25} (1 \pm 3.35 \times 10^{-4}) = (2.20086 \pm 0.00074) \times 10^{25} \text{ cm}^3$$

77. Mean density:

$$\bar{\rho}_e = 3.3420 (1 \pm 1.5 \times 10^{-3}) = 3.3420 \pm 0.0050 \text{ g/cm}^3$$

78. Surface density:

$$\rho_0 = 3.290 \text{ g/cm}^3$$

79. Central density:

$$\rho_c = 3.420 \text{ g/cm}^3$$

80. Inhomogeneity factor of the Moon:

$$\lambda = 0.1991 (1 \pm 5.0 \times 10^{-4}) = 0.1991 \pm 0.0001$$

81. Dimensionless moment of inertia parameters:

$$f = \frac{C - B}{C - A} = 0.70 (1 \pm 2.86 \times 10^{-2}) = 0.70 \pm 0.02$$

$$\alpha = \frac{C - B}{C} = 0.0004395 (1 \pm 3.03 \times 10^{-2}) = 0.0004395 \pm 0.0000133$$

$$\beta = \frac{C - A}{C} = 0.0006279 (1 \pm 1.6 \times 10^{-4}) = 0.0006279 \pm 0.0000010$$

$$\gamma = \frac{B - A}{C} = 0.0001884 (1 \pm 6.85 \times 10^{-2}) = 0.0001884 \pm 0.0000129$$

$$g = \frac{3}{2} \frac{C}{M_e a^2} = 0.5972 (1 \pm 5.0 \times 10^{-4}) = 0.5972 \pm 0.0003$$

$$J = \frac{3}{2} \frac{C - 1/2 (A + B)}{M_e a^2} = 0.0003187 (1 \pm 1.38 \times 10^{-2}) = 0.0003187 \pm 0.0000044$$

$$K = \frac{3}{2} \frac{B - A}{M_e a^2} = 0.0001125 (1 \pm 6.84 \times 10^{-2}) = 0.0001125 \pm 0.0000077$$

$$L = \frac{3}{2} \frac{C - A}{M_e a^2} = 0.0003750 (1 \pm 2.1 \times 10^{-3}) = 0.0003750 \pm 0.0000008$$

82. Dimensionless moment of inertia differences:

$$\frac{C - A}{M_e a^2} = 0.0002500 (1 \pm 2.1 \times 10^{-3}) = 0.0002500 \pm 0.0000005$$

$$\frac{C - B}{M_e a^2} = 0.0001750 (1 \pm 3.09 \times 10^{-2}) = 0.0001750 \pm 0.0000054$$

82. Dimensionless moment of inertia differences (Continued):

$$\frac{B - A}{M_{\bullet} a^2} = 0.0000750 (1 \pm 6.84 \times 10^{-2}) = 0.0000750 \pm 0.0000051$$

83. Dimensionless moments of inertia:

$$\frac{A}{M_{\bullet} a^2} = 0.39787_7 (1 \pm 5.0 \times 10^{-4}) = 0.39787_7 \pm 0.00020$$

$$\frac{B}{M_{\bullet} a^2} = 0.39795_2 (1 \pm 5.0 \times 10^{-4}) = 0.39795_2 \pm 0.00020$$

$$\frac{C}{M_{\bullet} a^2} = 0.39812_7 (1 \pm 5.0 \times 10^{-4}) = 0.39812_7 \pm 0.00020$$

84. Axial ratios of the Moon:

$$\frac{b}{a} = \sqrt{1 - \frac{2\gamma}{1 + \gamma}} = 0.9998116 (1 \pm 1.3 \times 10^{-5}) = 0.9998116 \pm 0.0000129$$

$$\frac{c}{a} = \sqrt{1 - \frac{2\beta}{1 + \gamma}} = 0.9993720 (1 \pm 1.0 \times 10^{-6}) = 0.9993720 \pm 0.0000010$$

$$\sigma = R_{\bullet}/a = \frac{1}{2} \left( \frac{b}{a} + \frac{c}{a} \right) = 0.9995918 (1 \pm 7.0 \times 10^{-6}) = 0.9995918 \pm 0.0000070$$

85. Oblateness coefficients of the potential function of the Moon:

$$J_2 = \frac{C - 1/2(A + B)}{M_{\bullet} a^2} = \frac{2}{3} J = 0.0002125 \pm 0.0000029$$

$$J_2^{(2)} = \frac{B - A}{4 M_{\bullet} a^2} = \frac{1}{6} K = 0.0000188 \pm 0.0000013$$

86. Unit for the Moon's moments of inertia:

$$M_{\bullet} a^2 = 2.2241_6 \times 10^{35} (1 \pm 1.2 \times 10^{-3}) = (2.2241_6 \pm 0.0027) \times 10^{35} \text{ kg m}^2$$

87. Moment of inertia differences:

$$C - \frac{1}{2}(A + B) = 0.000473 \times 10^{35} (1 \pm 1.50 \times 10^{-2}) = (0.000473 \pm 0.000007) \times 10^{35} \text{ kg m}^2$$

$$B - A = 0.000167 \times 10^{35} (1 \pm 6.96 \times 10^{-2}) = (0.000167 \pm 0.000012) \times 10^{35} \text{ kg m}^2$$

88. Moon's moments of inertia:

$$A = 0.88494 \times 10^{35} (1 \pm 1.73 \times 10^{-3}) = (0.88494 \pm 0.00153) \times 10^{35} \text{ kg m}^2$$

$$B = 0.88511 \times 10^{35} (1 \pm 1.73 \times 10^{-3}) = (0.88511 \pm 0.00153) \times 10^{35} \text{ kg m}^2$$

$$C = 0.88550 \times 10^{35} (1 \pm 1.73 \times 10^{-3}) = (0.88550 \pm 0.00153) \times 10^{35} \text{ kg m}^2$$

## 1. THE ASTRONOMICAL UNIT AND THE SOLAR PARALLAX

The astronomical unit (A.U.), or the Earth's mean distance from the Sun, is connected with the solar parallax ( $\pi_\odot$ ) by the following relation (with  $R_\oplus = 6378.170 \pm 0.020$  km the equatorial radius of the Earth):

$$1 \text{ A.U.} = \frac{R_\oplus}{\sin \pi_\odot} = \frac{R_\oplus}{\pi_\odot'' \sin 1''} = \frac{206264.''806247}{\pi_\odot''} R_\oplus = \frac{1315592000 \pm 4000}{\pi_\odot''} \quad (1)$$

Modern determinations of the solar parallax usually are included between the two values  $\pi_\odot = 8.''790 \pm 0.''001$  (H. Spencer Jones, 1941) and  $\pi_\odot = 8.''79835 \pm 0.''00039$  (E. Rabe, 1949). The mean value of both determinations,  $\pi_\odot = 8.''794 \pm 0.''002$ , has been accepted by C. W. Allen (Ref. 1, p. 131) in his book on "Astrophysical Quantities" (1955). Exactly in the middle of these two values are also the recently obtained data of radar echoes from Venus, which have a considerably higher accuracy than previous determinations. Furthermore, agreement of the different radio observatories is also very good, as shown in the following table:

Radio Observatory	Author (Year)	Ref.	Radar Frequency (MC/sec)	Astronomical Unit	Solar Parallax ( $R_\oplus = 6378.170$ km)
Millstone (Lincoln Lab., M.I.T.)	Pettengill, Price et. al. (1961)	2	440	$149597850 \pm 400$	$8.''79419_1$
Goldstone (J.P.L.)	Victor, Stevens and Muhlemann (1961)	3	2388	$149598845 \pm 250$	$8.''79413_2$
Jodrell Bank (U. of Manchester)	Thomson et. al. (1961)		408	$149601000 \pm 5000$	$8.''79400_5$
Moorestown (R.C.A.)	Maron et. al. (1961)		---	$149596000$	$8.''79429_9$
U. S. S. R.	Kotelnikov (1961)		700	$149599500 \pm 800$	$8.''79409_4$
					$8.''79414_4$

According to Newcomb and de Sitter the semimajor axis of the Earth's orbit around the Sun is given by  $a_\oplus = 1.000000236 + 0.000000004$  A.U. (or approximately 35 km more than 1 A.U.). For practical purposes both distances will be assumed equal to

$$a_\oplus = 149598700 \pm 400 = 149598700 (1 \pm 2.7 \times 10^{-6}) \text{ km} \quad (2)$$

The corresponding solar parallax will be

$$\pi_\odot = \frac{R_\oplus}{a_\oplus \sin 1''} = 206264.''806247 \frac{6378.170 \pm 0.020}{149598700 \pm 400} = 8.''79414 \pm 0.''00005 \quad (3)$$

Taking as best value for the light velocity the value determined by Froome (Ref. 4) in 1958:

$$c = 299792.5 \pm 0.1 = 299792.5 (1 \pm 3.3 \times 10^{-7}) \text{ km/sec} \quad (4)$$

the light-time for unit distance (1 A.U.) is therefore

$$r = \frac{a_{\oplus}}{c} = 499.007_s \pm 0.018 = 499.007_s (1 \pm 3.6 \times 10^{-5}) \text{ sec} \quad (5)$$

## 2. DEFINITION OF TIME UNITS, MEAN ORBITAL MOTIONS, AND ROTATIONAL ANGULAR VELOCITIES OF EARTH AND MOON

There are three different times which are in use, namely, the Greenwich mean solar time or universal time (U.T.), the Greenwich mean sidereal time (G.M.S.T.) and the ephemeris time (E.T.) or Newtonian time. Due to the variable rotation of the Earth, the mean solar time and the mean sidereal time do not have a constant rate. The observations are therefore functions of a variable time, while the gravitational theories for the Sun and the planets use a uniform time. The ephemeris time, having a constant rate is defined by the orbital motion of the Earth as given by Newcomb's Tables of the Sun. It is therefore necessary to apply corrections to our practical determinations of time. In addition to the fluctuations and the tidal slowing down of the Earth's rotation, the Moon also shows a real diminution in the angular mean motion which is not given by Brown's lunar theory.

The correction to Newcomb's tabulated tropical mean longitude of the Sun (Ref. 5)

$$L_{\odot} = 279^{\circ} 41' 48.''04 + 129602768.''13 T_E + 1.''089 T_E^2 \quad (6)$$

is, according to H. Spencer Jones (Ref. 6),

$$\Delta L_{\odot} = +1.''00 + 2.''97 T + 1.''23 T^2 + 0.0748 B \quad (7)$$

when the observation times are in U.T. The time  $T$  is in Julian centuries of  $36525^d$  counted from 1900 Jan. 0, 12<sup>h</sup> U.T. (Greenwich mean noon) and  $B$  is the irregular fluctuation in the Moon's mean longitude in arc seconds (time of observation again expressed in U.T.). The Sun's tropical mean longitude,  $L_{\odot}$ , increases at the rate of  $1''$  in  $86400/(0.9856473354 \times 3600) = 24.34948$  sec, so that the correction to universal time, required to obtain ephemeris time is, according to H. Clemence (Ref. 7),

$$\Delta t \equiv t_E - t_U = 24.34948 \Delta L_{\odot} = +24^5.349 + 72^5.318 T + 29^5.950 T^2 + 1.82134 B \quad (8)$$

H. Spencer Jones gives for the irregular fluctuation (Ref. 6)

$$B = (L_{\odot, \text{obs.}} - L_{\odot, \text{tabular}}) + 10.''71 \sin(140^{\circ}0' T + 240^{\circ}7') - 4.''65 - 12.''96 T - 5.''22 T^2 \quad (9)$$

The periodic term is Brown's empirical term in his lunar theory. Therefore the correction to the Moon's mean longitude, as given by Brown's Tables of the Motion of the Moon (Ref. 8), is

$$\Delta L_{\oplus} \equiv L_{\oplus, \text{obs.}} - L_{\oplus, \text{tabular}} = +4.''65 + 12.''96 T + 5.''22 T^2 + B - 10.''71 \sin(140^{\circ}0' T + 240^{\circ}7') \quad (10)$$

in order to obtain the actual mean longitude determined by observations in U.T. In the time

interval  $\Delta t = t_E - t_U$  the Moon's mean longitude increases by ( $n = 13.176\ 396\ 5268 \times 3600/86400 = 0.549\ 016\ 522''/\text{sec}$ )

$$\Delta L_t = 0.549 \Delta t = 13.368 \Delta L_\odot = +13.''37 + 39.''70 T + 16.''44 T^2 + B \quad (11)$$

Therefore the correction to Brown's Tables is

$$\Delta L_t - \Delta L_\odot = -8.''72 - 26.''74 T - 11.''22 T^2 - 10.''71 \sin(140^\circ 0' T + 240^\circ 7') \quad (12)$$

when the observations are in ephemeris time. Brown's theory is now reduced to a gravitational theory with the same measure of time as defined by Newcomb's Tables of the Sun. Clemence's corrected value for the Moon's mean longitude (Ref. 7)

$$L_t = 270^\circ 26' 2.''99 + 1732\ 564\ 379.''31 T_E - 4.''08 T_E^2 + 0.''0068 T_E^3 \quad (13)$$

is used in the American Ephemeris and Nautical Almanac.

By means of equation (6) the tropical year, from mean equinox to mean equinox, thus has the length

$$\begin{aligned} P_{\text{trop}} &= \frac{2\pi}{L_\odot} = \frac{1\ 296\ 000'' \times 36\ 525^{dE}}{129\ 602\ 768.''13 + 2.''178 T} = 365.^{dE}242\ 198\ 78 - 0.^{dE}000\ 006\ 138 T \\ &= 365^{dE}05^{hE} 48^{mE} 45.^s E 9747 - 0.^s E 5303 T = 31\ 556\ 925.^s E 9747 - 0.^s E 5303 T \end{aligned} \quad (14)$$

In 1957 the ephemeris second has been adopted as the fundamental invariable unit of time, and it is the fraction  $1/31\ 556\ 925.9747$  of the tropical year for 1900 Jan. 0, 12<sup>h</sup> E.T. (Ref. 9)

The basis for all civil time-keeping is the universal time which is non-uniform. In practical life, however, the difference between mean solar time and ephemeris time can be neglected because there is  $1^d \approx (1 \pm 10^{-8})^{dE}$ . To define universal time Newcomb introduced a fictitious mean Sun which moves with the same constant sidereal rate, in the equator, as the mean sidereal motion is for the true Sun, affected by aberration (20.''50) in the ecliptic. According to Newcomb, the right ascension of the fictitious mean Sun is (neglecting nutation in right ascension)

$$\begin{aligned} R_E &= 279^\circ 41' 27.''54 + 129\ 602\ 768.''13 T_E + 1.''394 T_E^2 \\ &= 18^h 38^m 45.^s 836 + 8\ 640\ 184.^s 542 T_E + 0.^s 0929 T_E^2 \end{aligned} \quad (15)$$

Defining a point on the equator whose right ascension, measured from the mean equinox of date, is

$$R_U = 18^h 38^m 45.^s 836 + 8\ 640\ 184.^s 542 T_U + 0.^s 0929 T_U^2, \quad (16)$$

and where  $R_E$  differs from  $R_U$  by 0.002738  $\Delta t$  (see equation 8), the Greenwich hour angle

$\tau_{Gr}(R_U)$  of the point whose right ascension is  $R_U$  [equal to universal time (U.T.)  $\pm 12^h$ ], increased by the right ascension  $R_U$ , is the Greenwich hour angle,  $\tau_{Gr}(\gamma)$ , of the mean vernal equinox of date which is Greenwich mean sidereal time,  $\Theta_{Gr}$ . That is,

$$U.T. \pm 12^h + R_U = \tau_{Gr}(R_U) + R_U = \tau_{Gr}(\gamma) \equiv \Theta_{Gr} \quad (17)$$

Adding the East longitude to both sides gives the local mean time on the left side of the equation and the local mean sidereal time on the right side, because

$$\Theta = \Theta_{Gr} + \lambda_{east}.$$

The time rate of the right ascension, given by differentiation of equation (16), is

$$\begin{aligned} \dot{R}_U &= 8640184.542 + 0.1858 T_U \text{ [sec/Jul. century]} \\ &= 129602768.13 + 2.788 T_U \text{ ["/Jul.century]} \\ &= 3548.3304074 + 0.00007633 T_U \text{ ["/d]} \\ &= 0.9856473354 + 2.1203 \times 10^{-8} T_U \text{ [°/d]}, \end{aligned} \quad (18)$$

Adding to this the time rate of the hour angle,  $\dot{\gamma} = 360 \text{ °/d} = 1296000 \text{ "/d}$ , the time rate of the mean sidereal time is then

$$\begin{aligned} \dot{\Theta}_m &= 1299548.3304074 + 0.00007633 T \text{ ["/d]} \\ &= 360.9856473354 + 2.1203 \times 10^{-8} T \text{ [°/d]} \\ &= 15.04106863897 + 8.835 \times 10^{-10} T \text{ ["/s or °/h]} \\ &= 1.002737909265 + 0.5890 \times 10^{-10} T \text{ [d*/d or s*/s]} \\ &= 7.29211585458 \times 10^{-5} + 4.283 \times 10^{-15} T \text{ [rad/s]} \end{aligned} \quad (19)$$

This motion is the result of the spin of the Earth and the motion of the vernal equinox (precession). Because the latter motion takes place in the ecliptic the equatorial component of the general precession in right ascension,  $m$ , must be used here. The mean angular velocity of the Earth's rotation is, therefore,

$$\Omega_m = \dot{\Theta}_m - m \quad (20)$$

It is very probable that Newcomb's value for the general precession in longitude,  $p$ , must be increased by  $\Delta p = + 0.^{\circ}80$  per tropical century (see Part 9), thus

$$\begin{aligned} p &= 5026.441 + 2.2229 T + 0.00026 T^2 \text{ ["/trop. century]} \\ &= 0.1376194 + 0.000060861 T + 0.712 \times 10^{-8} T^2 \text{ ["/d]} \\ &= 0.00003822761 + 0.000000016906 T + 1.98 \times 10^{-12} T^2 \text{ [°/d]} \end{aligned} \quad (21)$$

Neglecting the correction in planetary precession ( $\Delta\lambda = 0$ ) the correction for the general precession in right ascension would be  $\Delta m = \Delta p \cdot \cos \epsilon = 0.^s 80 \times 0.917 = + 0.^s 73$ . Thus,

$$\begin{aligned} m &= 4609.236 + 2.7945 T + 0.00012 T^2 [''/\text{trop. century}] \\ &= 0.1261967 + 0.000076511 T + 0.33 \times 10^{-8} T^2 [''/\text{d}] \\ &= 0.00003505464 + 0.000000021253 T + 0.92 \times 10^{-12} T^2 [\text{°}/\text{d}] \end{aligned} \quad (22)$$

The angular velocity of the Earth's rotation is, therefore,

$$\begin{aligned} \Omega_m &= 1299548.2042107 - 0.00000018 T [''/\text{d}] \\ &= 360.9856122808 - 0.0050 \times 10^{-8} T [\text{°}/\text{d}] \\ &= 15.04106717837 - 0.021 \times 10^{-10} T [''/\text{s} \text{ or } \text{°}/\text{h}] \\ &= 1.002737811891 - 0.0014 \times 10^{-10} T [\text{rot}/\text{d}] \\ &= 7.29211514646 \times 10^{-5} - 0.010 \times 10^{-15} T [\text{rad}/\text{s}] \end{aligned} \quad (23)$$

Using equations (19) and (23) the following periods are obtained:

1. Mean solar day (culmination period of the mean Sun)

$$\begin{aligned} 1^d &= 1^{d*} 002737909265 + 0^{d*} 589 \times 10^{-10} T \\ &= 86636.^s 55536050 + 0.^s 0508896 \times 10^{-4} T \\ &= 24^h 03^m 56.^s 55536050 + 0.^s 0508896 \times 10^{-4} T \\ &= 1.^{\text{rot}} 002737811891 - 0.^{\text{rot}} 0014 \times 10^{-10} T \\ &= (1 + 10^{-8})^{dE} \end{aligned} \quad (24)$$

2. Mean siderial day or mean equinoctial day (culmination period of the vernal equinox)

$$\begin{aligned} 1^{d*} &= 1^d - 0.^d 002730433586 - 0.^d 587 \times 10^{-10} T \\ &= 0.^d 997269566414 - 0.^d 587 \times 10^{-10} T \\ &= 86164.^s 09053817 - 0.^s 0507168 \times 10^{-4} T \\ &= 23^h 56^m 04.^s 09053817 - 0.^s 0507168 \times 10^{-4} T \\ &= \frac{\Omega}{\Theta} = 1 - \frac{m}{\Theta} = (1 - 0.000000097108)^{\text{rot}} - 0.^{\text{rot}} 589 \times 10^{-10} T \\ &= 0.^{\text{rot}} 99999902892 - 0.^{\text{rot}} 589 \times 10^{-10} T \end{aligned} \quad (25)$$

3. Period of the Earth's rotation (culmination period of an equatorial star without proper motion)

$$\begin{aligned}
 \dot{\Omega}_{\text{rot}} &= \frac{\dot{\Theta}}{\Omega} = 1 + \frac{m}{\Omega} = 1^d 000 000 097 108 + 0^d 589 \times 10^{-10} T \\
 &= 86400^s 008 390 13 + 0^s 050 889 6 \times 10^{-4} T \\
 &= 1^d - 0^d 002 730 336 743 + 0^d 001 4 \times 10^{-10} T \\
 &= 1^d - 235^s 901 094 60 + 0^s 000 120 96 \times 10^{-4} T \\
 &= 0^d 997 269 663 257 + 0^d 001 4 \times 10^{-10} T \\
 &= 86164^s 098 905 40 + 0^s 001 209 6 \times 10^{-5} T \quad (26)
 \end{aligned}$$

Because

$$\begin{aligned}
 1^d - 1^d * &= 236^s 555 360 50 + 0^s 050 889 6 \times 10^{-4} T \\
 &= 235^s 909 461 83 + 0^s 050 716 8 \times 10^{-4} T
 \end{aligned}$$

or

$$1^h - 1^h * = 9^s 856 473 = 9^s 829 561$$

the change of mean siderial time against mean solar time is  $9^s 856 47$  in a mean solar hour and  $9^s 829 56$  in a mean siderial hour.

In order to apply Kepler's third law, the siderial mean angular motions of the Earth about the Sun and the Moon about the Earth will be needed. Differentiation of equation (6) gives the tropical mean motion of the Earth:

$$\begin{aligned}
 n_{\oplus} (\text{trop.}) &= \dot{L}_{\oplus} = 129 602 768.13 + 2.178 T [\text{''/Jul. century}] \\
 &= 3548.330 407 4 + 0.000 059 63 T [\text{''/d}] \\
 &= 0.985 647 335 4 + 1.656 4 \times 10^{-8} T [\text{°/d}] \quad (27)
 \end{aligned}$$

Subtracting from this the general precession in longitude (equation 21) yields the siderial mean motion of the Earth:

$$\begin{aligned}
 n_{\oplus} (\text{sid.}) &= \dot{L}_{\oplus} - p = 3548.192 788 0 - 0.000 001 23 T [\text{''/d}] \\
 &= 0.985 609 107 8 - 0.034 2 \times 10^{-8} T [\text{°/d}] \\
 &= 0.041 067 046 16 - 0.142 4 \times 10^{-10} T [\text{''/s}] \\
 &= 1.990 986 582 0 \times 10^{-7} - 0.690 374 68 \times 10^{-16} T [\text{rad/s}] \quad (28)
 \end{aligned}$$

Differentiating equation (13) provides the tropical mean motion of the Moon:

$$\begin{aligned}
n_4 (\text{trop}) &= \dot{L}_4 = 1732564379.31 - 8.16 T + 0.0204 T^2 [\text{"/Jul. century}] \\
&= 47435.0274965 - 2.234 \times 10^{-4} T + 0.559 \times 10^{-6} T^2 [\text{"/d}] \\
&= 13.1763965268 - 6.206 \times 10^{-8} T + 1.55 \times 10^{-10} T^2 [\text{"/d}] \\
&= 0.549016521950 - 2.586 \times 10^{-9} T + 6.46 \times 10^{-12} T^2 [\text{"/s}] \quad (29)
\end{aligned}$$

and subtraction of the general precession in longitude,  $p$ , gives the siderial mean motion of the Moon:

$$\begin{aligned}
n_4 (\text{sid.}) &= \dot{L}_4 - p = 47434.8898771 - 2.843 \times 10^{-4} T + 0.552 \times 10^{-6} T^2 [\text{"/d}] \\
&= 13.1763582992 - 7.897 \times 10^{-8} T + 1.53 \times 10^{-10} T^2 [\text{"/d}] \\
&= 0.549014929133 - 3.290 \times 10^{-9} T + 6.38 \times 10^{-12} T^2 [\text{"/s}] \\
&= 2.66169948773 \times 10^{-6} - 1.595 \times 10^{-13} T + 3.09 \times 10^{-17} T^2 [\text{rad/s}] \quad (30)
\end{aligned}$$

Because the distance of the Earth to the Sun is now known more accurately than before, it is possible to give the mean orbital velocity of the Earth about the Sun with high accuracy, namely

$$v_{\oplus} = a_{\oplus} n_{\oplus} = 29784.90 \pm 0.08 [\text{m/s}] \quad (31)$$

There are two constants connected with this velocity. Taking  $e = 0.01675$  for the orbital eccentricity of the Earth the value for the constant of aberration will be

$$K = \frac{v_{\oplus}/c}{\sqrt{1-e^2} \sin 1''} = \frac{\tau n_{\oplus}}{\sqrt{1-e^2} \sin 1''} = 20.''4956 \pm 0.''0007, \quad (32)$$

and using the formula of de Sitter (Ref. 10) the geodetic precession, due to the special theory of relativity, is

$$p_g = \frac{3}{2} (v_{\oplus}/c)^2 n_{\oplus} = \frac{3}{2} (K \sin 1'' \sqrt{1-e^2})^2 n_{\oplus} = 1.''9188 \pm 0.''0002 \quad (33)$$

### 3. THE LUNAR DISTANCE AND THE LUNAR PARALLAX

The mean observed distance,  $\bar{r}_{\oplus}$ , of the Moon from the Earth is connected with the mean perturbed lunar parallax,  $\pi_4$ , and the constant  $\pi'_4$  of the sine of the perturbed lunar parallax by the following relation:

$$\frac{\bar{r}_{\oplus}}{R_{\oplus}} = \frac{1}{\sin \pi_4} = \frac{1}{\pi'_4 \sin 1''} \quad (34)$$

Dividing both sides of the series development

$$\pi_4 = \sin \pi_4 + \frac{1}{6} \sin^3 \pi_4 + \dots$$

by  $\sin 1''$  yields (because  $\pi'_e = \sin \pi_e / \sin 1''$ )

$$\begin{aligned}\pi''_e &= \pi'_e + \frac{1}{6} (\pi'_e)^3 (\sin 1'')^2 + \dots = \pi'_e \left[ 1 + \frac{1}{6} \left( \frac{R_e}{\bar{r}_e} \right)^2 + \dots \right] \\ &= \pi'_e \times 1.000045885 = \pi'_e + 0.''157\end{aligned}\quad (35)$$

Newer determinations are

$\pi'_e = 3422.''54$	$\pi_e = 3422.70$	(E.W. Brown, Ref. 8)
$3422.526 \pm 0.009$	$3422.683 \pm 0.009$	(W. deSitter, Ref. 10)
$3422.419 \pm 0.024$	$3422.576 \pm 0.024$	(H. Jeffreys, Ref. 11, p. 193)
$3422.493$	$3422.650$	(Herrick, Baker, Ref. 12)

Recent determinations of the mean lunar distance,  $\bar{r}_e$ , by means of radar echoes to the Moon are in very close agreement (see Reference 13). This value is given by

$$\bar{r}_e = 384402.0 + 1.0 = 384402.0 (1 \pm 2.6 \times 10^{-6}) \text{ km} \quad (36)$$

thus

$$\frac{\bar{r}_e}{R_e} = \frac{384402.0 (1 \pm 2.6 \times 10^{-6})}{6378.170 (1 \pm 3.2 \times 10^{-6})} = 60.26838 (1 \pm 5.8 \times 10^{-6}) = 60.26838 \pm 0.00035 \quad (37)$$

and

$$\pi'_e = \frac{\sin \pi_e}{\sin 1''} = \frac{206264.''806247}{(\bar{r}_e/R_e)} = 3422.''438 \pm 0.''020 \quad (38)$$

$$\pi_e = 3422.''595 + 0.''020 \quad (39)$$

To obtain the semi-major axis,  $a_e$ , it is necessary to add to the mean lunar distance the constant part of the solar perturbations according to Brown's lunar theory. There is now

$$a_e = 1.000907681 \bar{r}_e = 384750.9 \pm 1.0 \text{ km} \quad (40)$$

and the mean orbital velocity of the Moon about the Earth is

$$v_e = a_e \pi_e = 1024.091 \pm 0.003 \text{ m/s} \quad (41)$$

#### 4. MASS RATIOS OF THE SUN AND THE EARTH-MOON SYSTEM

Taking the already given values for  $a_\oplus$ ,  $n_\oplus$  (sid.), and  $a_e$ ,  $n_e$  (sid.) then Kepler's third law gives

$$n_\oplus^2 a_\oplus^3 = G (M_\odot + M_\oplus + M_e) = G M_\oplus (\nu + 1)(1 + \kappa) \quad (42)$$

$$n_e^2 a_e^3 = G (M_\oplus + M_e) = G M_\oplus (1 + \kappa) \quad (43)$$

with

$$\nu = \frac{M_\oplus}{M_\oplus + M_\star} \gg 1 \quad ; \quad \kappa = \frac{M_\star}{M_\oplus} \ll 1 \quad (44)$$

Dividing the two equations yields

$$\begin{aligned} \nu &= \frac{M_\oplus}{M_\oplus + M_\star} = \left(\frac{n_\oplus}{n_\star}\right)^2 \left(\frac{a_\oplus}{a_\star}\right)^3 - 1 = \left(\frac{\nu_\oplus}{\nu_\star}\right)^2 \left(\frac{a_\oplus}{a_\star}\right)^3 - 1 \\ &= 328898.6 (1 \pm 1.6 \times 10^{-5}) = 328898.6 \pm 5.2 \end{aligned} \quad (45)$$

This value is approximately in the middle between the value 329390 obtained by S. Newcomb (Ref. 14), adopted by Am. Ephemeris, and the value 328446 ± 43 determined by E. Rabe (Ref. 15).

The second equation gives

$$\mu_\oplus (1 + \kappa) = n_\star^2 a_\star^3 = \nu_\star^2 a_\star = 403512.3 \pm 3.2 \text{ km}^3/\text{s}^2 \quad (46)$$

which connects the gravitational parameter  $\mu_\oplus = GM_\oplus$  for the Earth with  $\kappa$ , the ratio of the Moon's mass to the Earth's mass.

## 5. THE CONSTANT OF LUNAR INEQUALITY AND THE PARALLACTIC INEQUALITY IN MOON'S ECLIPTIC LONGITUDE

The Parallactic Inequality in the Moon's ecliptic longitude is given by E. W. Brown's lunar theory as follows

$$P_* = (49853.''2 \pm 1.''2) \frac{1 - \kappa}{1 + \kappa} \frac{\pi_\oplus}{\pi_\star} = (49853.''2 \pm 1.''2) \frac{1/\kappa - 1}{1/\kappa + 1} \frac{\bar{r}_\star}{a_\oplus} \quad (47)$$

or with the newest data for the lunar distance and the astronomical unit

$$P_* = (128.''1005 \pm 0.''0037) \frac{1/\kappa - 1}{1/\kappa + 1} \quad (48)$$

Newer determinations are:

$$\begin{aligned} P_* &= 124.''86 \pm 0.''15 && (\text{J. Bauschinger, Ref. 16}) \\ &= 125.154 && (\text{E. W. Brown, Ref. 8}) \\ &= 124.93 && (\text{H. Battermann, Ref. 17}) \\ &= 124.969 \pm 0.042 && (\text{D. Brouwer and O. B. Watts, Ref. 18}) \end{aligned}$$

On the other hand, the constant of Lunar Inequality is defined by W. deSitter (Ref. 10) as

$$L = \frac{\kappa}{1 + \kappa} \frac{\pi_\oplus}{\sin \pi_\star} = \frac{206264.''806247}{1/\kappa + 1} \frac{\pi_\oplus}{\pi_\star} = \frac{530.''0089 \pm 0.''0028}{1/\kappa + 1} \quad (49)$$

Newcomb used the lunar inequality in the Sun's longitude which is, according to deSitter,  $L_s = 1.00450L$ . The ratio of  $P_*$  and  $L$ , depending only on  $\kappa$ , is

$$P_* / L = \frac{49.853.^{\prime\prime}2 \pm 1.^{\prime\prime}2}{206.264.^{\prime\prime}806.247} (1/\kappa - 1) = (0.241695 \pm 0.000006) (1/\kappa - 1) \quad (50)$$

The mass ratio is therefore

$$1/\kappa = (4.13744_4 \pm 0.00010) \frac{P_*}{L} + 1 = \frac{530.^{\prime\prime}0089 \pm 0.^{\prime\prime}0028}{L} - 1 = \frac{128.^{\prime\prime}1005 + P_*}{128.^{\prime\prime}1005 - P_*} \quad (51)$$

Observed values for the constant of Lunar Inequality are:

$L = 6.^{\prime\prime}456 \pm 0.^{\prime\prime}012$	(Newcomb, Ref. 14)	From observations of	Sun
$6.414 \pm 0.009$	(D. Gill, Ref. 19)	"	Victoria
$6.4305 \pm 0.0031$	(A.R. Hinks, Ref. 20)	"	Eros (opp. of 1901)
$6.4390 \pm 0.0015$	(H. Spencer Jones, Ref. 21)	"	Eros (opp. of 1931)
$6.450 \pm 0.010$	(Morgan and Scott, Ref. 22)	"	Sun
$6.4378 \pm 0.0015$	(H. Jeffreys, Ref. 23)	"	Eros (opp. of 1931)
$6.4356 \pm 0.0028$	(E. Rabe, Ref. 15)	"	Eros (opp. of 1931)
$6.4428 \pm 0.0014$	(E. Delano, Ref. 24)	"	Eros (opp. of 1931)
$6.4430 \pm 0.0017$			

The latest reevaluation of all Eros observations during the opposition of 1930/31 by E. Delano (Ref. 24) gave

$$\begin{aligned} L &= 6.^{\prime\prime}4428 \pm 0.^{\prime\prime}0014 && (\text{from right ascensions of Eros}) \\ L &= 6.4430 \pm 0.0017 && (\text{from declinations of Eros}) \end{aligned}$$

Delano used the old value  $\pi_0 = 8.^{\prime\prime}790$  for the solar parallax and obtained therefore  $1/\kappa = 81.222 \pm 0.027$  and  $81.219 \pm 0.030$ , respectively. With the newest values for  $\pi_0$  and  $\pi'$  there is now

$$1/\kappa = 81.263_7 \quad \text{and} \quad 1/\kappa = 81.261_2$$

respectively.

## 6. ANOTHER METHOD FOR THE DETERMINATION OF THE RATIO OF THE MASSES OF EARTH AND MOON

The mass of the Earth is given by

$$M_\oplus = \frac{4}{3} \pi R_\oplus^3 (1-f) \rho_\oplus \quad (52)$$

where  $f$  is the flattening (oblateness) and  $\rho_\oplus$  is the mean density of the Earth. On the other hand the mass of the triaxial figure of the Moon is given by

$$M_\bullet = \frac{4}{3} \pi a b c \rho_\bullet = \frac{4}{3} \pi a^3 \left( \frac{b}{a} \right) \left( \frac{c}{a} \right) \rho_\bullet \quad (53)$$

Because the longest axis is always directed to the Earth (neglecting the small librations) and can never be seen, the lunar radius of the visible disk is

$$R_{\text{e}} = \frac{b+c}{2} ; \quad \frac{R_{\text{e}}}{a} = \frac{1}{2} \left( \frac{b}{a} + \frac{c}{a} \right) = \sigma \quad (54)$$

That yields for the mass of the moon

$$M_{\text{e}} = \frac{4}{3} \pi R_{\text{e}}^3 \frac{(b/a)(c/a)}{\sigma^3} \rho_{\text{e}} \quad (55)$$

The mass ratio is therefore

$$1/\kappa \equiv M_{\oplus}/M_{\text{e}} = \left( \frac{R_{\oplus}}{R_{\text{e}}} \right)^3 \frac{(1-f)\sigma^3}{(b/a)(c/a)} \left( \frac{\rho_{\oplus}}{\rho_{\text{e}}} \right) \frac{(1-f)\sigma^3}{(b/a)(c/a)} \frac{\rho_{\oplus}/\rho_{\text{e}}}{k^3} \quad (56)$$

where

$$k = \frac{R_{\text{e}}}{R_{\oplus}} = \frac{\sin s_{\text{e}}}{\sin \pi_{\text{e}}} = \frac{s_{\text{e}}'}{\pi_{\text{e}}} = \frac{s_{\text{e}} - 0.^{\circ}003}{\pi_{\text{e}} - 0.^{\circ}157} \quad (57)$$

is given by the lunar parallax,  $\pi_{\text{e}}$ , and the apparent semi-diameter of the Moon,  $s_{\text{e}}$ .

A reevaluation of Sir Harold Jeffrey's best data on the Moon's figure by the author gave (see section 12)

$$\frac{b}{a} = 0.9998116 ; \quad \frac{c}{a} = 0.9993720 ; \quad \sigma = 0.9995918$$

From the secular perturbations of artificial Earth satellites there follows as best value for the Earth oblateness

$$1/f = 298.30 ; \quad 1-f = 0.99664767$$

so that

$$1/\kappa = 0.9962409 \frac{\rho_{\oplus}/\rho_{\text{e}}}{k^3} \quad (58)$$

Taking for the mean densities the well-known and frequently used values

$$\rho_{\oplus} = 5.517 \pm 0.004 \text{ g/cm}^3 \text{ (Heyl)} ; \quad \rho_{\text{e}} = 3.342 \pm 0.005 \text{ g/cm}^3 \text{ (Jeffreys)}$$

the density ratio, independent from the assumed value of the gravitational constant,  $G$ , becomes  $\rho_{\oplus}/\rho_{\text{e}} = 1.6508$ . Therefore

$$1/\kappa = 1.6446/k^3 \quad (59)$$

With  $\pi' = 3422.''438$  and various values for  $s_e$  the following table is obtained

$s_e$	$k$	$1/\kappa$
932.''58 (Newcomb, Ref. 25)	0.2724891	81.286
932.63 (American Ephem., Ref. 26)	0.2725037	81.272
932.80 + 0.07 (Hirose & Manabe, Ref. 27)	0.2725534	81.228

The American Ephemeris is using  $k = 0.2724953$ , based on Brown's lunar parallax. The values for  $1/\kappa$  obtained in the previous paragraph are between the two latter values in this table. The arithmetic mean of these two latter values will be taken as the presently best value, namely

$$1/\kappa \equiv \frac{M_\oplus}{M_e} = 81.250 (1 \pm 3 \times 10^{-4}) = 81.250 \pm 0.024 \quad (60)$$

and therefore

$$k \equiv \frac{R_e}{R_\oplus} = 0.2725289 \pm 0.0000273 \quad (61)$$

The last equation of paragraph 3 now gives

$$\mu_\oplus \equiv GM_\oplus = \frac{403512.3}{1.0123077} = 398606.4 \pm 4.9 \text{ km}^3/\text{sec}^2 \quad (62)$$

for the Earth, while, for the Moon,

$$\mu_e \equiv GM_e = \kappa\mu_\oplus = 4905.92 \pm 1.52 \text{ km}^3/\text{sec}^2 \quad (63)$$

## 7. GEODYNAMIC (TERRESTRIAL) RELATIONS

The surface of the Earth (geoid) can be approximated as the surface of an spheroid assumed as an equipotential surface. The equation for the Earth's radius, as function of the latitude, is then given by

$$R = R_e \left[ 1 - f \sin^2 \varphi + \left( \frac{5}{8} f^2 - \kappa \right) \sin^2 2\varphi \right] = R_e \left[ 1 - f \sin^2 \phi - \left( \frac{3}{8} f^2 + \kappa \right) \sin^2 2\phi \right] \quad (64)$$

where  $\varphi$  is the geodetic (geographic) latitude and  $\phi$  the geocentric latitude. They are related by

$$\tan \phi = (1 - f)^2 \tan \varphi = (1 - e^2) \tan \varphi \quad (65)$$

where  $e = \sqrt{f(2-f)}$  is the eccentricity of the meridian ellipse of the Earth. The equation of the Earth-ellipsoid is obtained by setting  $\kappa = 0$ . The maximum depression,  $-\kappa R_e$ , of the spheroid from the ellipsoid is reached at the latitude  $45^\circ$ . It will never be more than 5.17 m. For the spheroid as equipotential surface there is

$$\begin{aligned} U &= \frac{GM}{R} \left[ 1 - \frac{2}{3} J \left( \frac{R}{R_e} \right)^2 P_2 (\sin \phi) + \frac{4}{15} K \left( \frac{R}{R_e} \right)^4 P_4 (\sin \phi) + \dots \right] + \frac{1}{2} \Omega^2 R^2 \cos^2 \phi \\ &= \frac{GM}{R} \left[ 1 - \sum_{n=2}^{\infty} J_n \left( \frac{R}{R_e} \right)^n P_n (\sin \phi) \right] + \frac{1}{2} \Omega^2 R^2 \cos^2 \phi = \text{const.} \end{aligned} \quad (66)$$

and the acceleration of gravity at this surface is

$$g = |\text{grad } U| = g_e [1 + \beta \sin^2 \varphi + \gamma \sin^2 2\varphi] = g_e [1 + \beta \sin^2 \phi + (\gamma + \beta f) \sin^2 2\phi] \quad (67)$$

where  $\Omega$  is the angular velocity of the Earth's rotation,  $\beta$  and  $\gamma$  are constant gravity coefficients and  $J_n$  or  $J$  and  $K$  are constant oblateness coefficients. These coefficients depend only on  $f$ ,  $\kappa$ , and a parameter containing  $\Omega^2$  (centrifugal force parameter). This latter parameter is a little different in the various second-order theories which have been developed. Taking

$$\Omega = 7.292115146 \times 10^{-5} \text{ rad/sec}$$

$$1/f = 298.30 \quad ; \quad 1 - f = 0.996647670$$

$$G = 6.670 \times 10^{-8} \text{ cm}^3/(\text{g} \cdot \text{sec}^2) \quad ; \quad \rho_m = 5.517 \text{ g/cm}^3 \quad ; \quad G \rho_m = 3.679839 \times 10^{-7} \quad (68)$$

then the following parameters could be used

$$\tilde{\omega} = \frac{\Omega^2 R_e^3}{GM} = \frac{\Omega^2}{4/3 \pi G \rho_m (1-f)} = 0.003461369 \quad [\text{Herrick, (Ref.12)}] \quad (69)$$

$$\tilde{\omega}' = \frac{\Omega^2 R_v^3}{GM} = \frac{\Omega^2}{4/3 \pi G \rho_m} = \tilde{\omega}(1-f) = 0.003449766 \quad [\text{Jeffreys, Ref.11}] \quad (70)$$

$$\tilde{\omega}_1 = \frac{\Omega^2 R_1^3}{GM} = \tilde{\omega} \left(1 - f + 2f^2 - \frac{8}{3}\kappa\right) = \tilde{\omega}' \left(1 + 2f^2 - \frac{8}{3}\kappa\right) = 0.003449843 \quad [\text{deSitter, Ref.10}] \quad (71)$$

$$\begin{aligned} m &= \frac{\Omega^2 R_e}{g_e} = \frac{\tilde{\omega}}{1 - A - \frac{3}{2}\tilde{\omega} + f + f^2 - \frac{3}{7}f\tilde{\omega} + \frac{16}{7}\kappa} = \frac{\tilde{\omega}}{1 - A - \tilde{\omega} + J + 1/2K} \\ &= 0.003467730 \quad [\text{Darwin, Ref.28; Helmert, Ref.29}] \end{aligned} \quad (72)$$

where  $R_v = R_e (1 - f)^{1/3}$  is the radius for a sphere of same volume as the Earth and  $R_1 = R_e (1 - 1/3f + 5/9f^2 - 8/9\kappa)$  is the mean radius for which  $P_2(\sin \varphi) = 0$  or  $\varphi = \sin^{-1}\sqrt{1/3} = 35^\circ 15' 51.''8$ .  $A = 0.88 \times 10^{-6}$  is the mass of the Earth's atmosphere (expressed in mass of the Earth) which does not contribute to the surface gravity of the Earth. Different assumptions have been made for  $\kappa$ . Bullard (Ref.30) found  $10^6 \kappa = 0.68$ . This value was accepted later by Herrick, Baker, and Hilton (Ref.12). On the other hand, deSitter found values of only  $10^6 \kappa = 0.47$  to  $0.52$  and used the round mean value  $10^6 \kappa = 0.50$ . The theoretical limits are according to deSitter (Ref.31)

$$0 \leq \kappa \leq \frac{5}{16}f\tilde{\omega} - \frac{1}{4}f^2 = (3.62 - 2.81) \times 10^{-6} = 0.81 \times 10^{-6} \quad (73)$$

The different formula systems now yield

$$\begin{aligned}
J &= f - \frac{1}{2} \tilde{\omega} - \frac{1}{2} f^2 + \frac{9}{14} f \tilde{\omega} + \frac{4}{7} \kappa \\
&= f - \frac{1}{2} \tilde{\omega}' - \frac{1}{2} f^2 + \frac{1}{7} f \tilde{\omega}' + \frac{4}{7} \kappa \\
&= f - \frac{1}{2} \tilde{\omega}_1 - \frac{1}{2} f^2 + \frac{1}{7} f \tilde{\omega}_1 + \frac{4}{7} \kappa \\
&= f - \frac{1}{2} m - \frac{1}{2} f^2 + \frac{1}{7} f m + \frac{3}{4} m^2 + \frac{4}{7} \kappa
\end{aligned} \tag{74}$$

$$\begin{aligned}
K &= \frac{6}{7} D = 3 f^2 - \frac{15}{7} f \tilde{\omega} + \frac{24}{7} \kappa \\
&= 3 f^2 - \frac{15}{7} f \tilde{\omega}' + \frac{24}{7} \kappa \\
&= 3 f^2 - \frac{15}{7} f \tilde{\omega}_1 + \frac{24}{7} \kappa \\
&= 3 f^2 - \frac{15}{7} f m + \frac{24}{7} \kappa
\end{aligned} \tag{75}$$

$$\begin{aligned}
\beta &= \frac{5}{2} \tilde{\omega} - f - \frac{26}{7} f \tilde{\omega} + \frac{15}{4} \tilde{\omega}^2 + \frac{8}{7} \kappa \\
&= \frac{5}{2} \tilde{\omega}' - f - \frac{17}{14} f \tilde{\omega}' + \frac{15}{4} \tilde{\omega}'^2 + \frac{8}{7} \kappa \\
&= \frac{5}{2} \tilde{\omega}_1 - f - \frac{17}{14} f \tilde{\omega}_1 + \frac{15}{4} \tilde{\omega}_1^2 + \frac{8}{7} \kappa \\
&= \frac{5}{2} m - f - \frac{17}{14} f m + \frac{8}{7} \kappa
\end{aligned} \tag{76}$$

$$\begin{aligned}
\gamma &= \frac{1}{8} f^2 - \frac{5}{8} f \tilde{\omega} - 3 \kappa \\
&= \frac{1}{8} f^2 - \frac{5}{8} f \tilde{\omega}' - 3 \kappa \\
&= \frac{1}{8} f^2 - \frac{5}{8} f \tilde{\omega}_1 - 3 \kappa \\
&= \frac{1}{8} f^2 - \frac{5}{8} f m - 3 \kappa
\end{aligned} \tag{77}$$

$$\begin{aligned}
\chi \equiv g_e / (GM/R_e^2) &= 1 - A - \tilde{\omega} + J + \frac{1}{2} K \\
&= 1 - A - \frac{3}{2} \tilde{\omega} + f + f^2 - \frac{3}{7} f \tilde{\omega} + \frac{16}{7} \kappa \\
&= 1 - A - \frac{3}{2} \tilde{\omega}' + f + f^2 - \frac{27}{14} f \tilde{\omega}' + \frac{16}{7} \kappa \\
&= 1 - A - \frac{3}{2} \tilde{\omega}_1 + f + f^2 - \frac{27}{14} f \tilde{\omega}_1 + \frac{16}{7} \kappa \\
&= 1 - A - \frac{3}{2} m + f + f^2 - \frac{27}{14} fm + \frac{9}{4} m^2 + \frac{16}{7} \kappa
\end{aligned} \tag{78}$$

Using these equations and the above-given constants for the Earth then the following table is obtained with  $J_2 \equiv (C - A)/(MR_e^2) = \frac{2}{3} J$  and  $J_4 = -\frac{4}{15} K = -\frac{8}{35} D$ :

Coefficient	$\kappa = 0$	$\kappa = 0.50 \times 10^{-6}$	$\kappa = 0.68 \times 10^{-6}$
$10^6 J$	1623.48	1623.77	1623.87
$10^6 K$	8.85	10.56	11.18
$10^6 J_2$	1082.32	1082.51	1082.58
$10^6 J_4$	- 2.36	- 2.82	- 2.98
$10^6 \beta$	5302.92	5303.49	5303.70
$10^6 \gamma$	- 5.85	- 7.35	- 7.89
$\chi$	0.9981 6566	0.9981 6680	0.9981 6721

The numerical values for  $|J_4|$  are a little higher than the values derived from the observed secular perturbations of artificial satellites. Thus the data for  $\kappa = 0$  will be used here.

The gravitational parameter of the Earth is now given by

$$\mu_{\oplus} \equiv GM_{\oplus} = \frac{4}{3} \pi G \rho_{\oplus} R_e^3 (1 - f) = \frac{g_e R_e^2}{\chi} = 398\,606.4 \pm 4.9 \text{ km}^3/\text{sec}^2 \tag{79}$$

which corresponds to  $1/\kappa = 81.250$ . Taking, furthermore,  $\frac{4}{3} \pi = 4.188\,790\,204$ ;  $1 - f = 0.996\,647\,670$ ;  $\chi = 0.998\,165\,66$  that yields

$$g_e R_e^2 = \chi \mu_{\oplus} = 397\,875.2 \pm 4.9 \text{ km}^2/\text{sec}^2 = 3.978\,752 \times 10^{14} \text{ m}^3/\text{sec}^2 \tag{80}$$

and

$$\frac{g_e}{R_e} = \frac{4}{3} \pi (1 - f) \chi G \rho_{\oplus} = 4.167\,090\,090 G \rho_{\oplus} \tag{81}$$

or with  $G = 6.670 \times 10^{-8} \text{ cm}^3/\text{g} \cdot \text{sec}^2$

$$\rho_{\oplus} = \frac{(g_e/R_e) \times 10^7}{2.77945} \quad [\text{g/cm}^3] \quad (82)$$

## 8. EXPERIMENTAL VALUES OF THE GRAVITATIONAL ACCELERATION OF THE EARTH

The observed gravitational acceleration at the surface of the rotating Earth can be represented by the formula

$$g = g_e [1 + \beta \sin^2 \varphi + \gamma \sin^2 2\varphi + \delta \cos^2 \varphi \cos 2(\lambda - \lambda_0)] \quad (83)$$

where  $\lambda$  is the geographic longitude measured eastwards of Greenwich meridian. The first term corresponds to a sphere. The next two terms give the contribution due to the oblateness of the Earth spheroid, while the longitude term is due to the non-ellipticity of the equator when the Earth is assumed as a triaxial figure. The longitude,  $\lambda_0$ , gives the direction of the longest semi-axis of the equator.  $\delta$  is connected with the difference  $B - A$  of the equatorial moments of inertia or with the flattening,  $f_e$ , of the equator by the relation

$$\delta = \frac{9}{4} \frac{B - A}{M_{\oplus} R_e^2} = \frac{9}{4} \Lambda f_e (2 - f_e) \approx \frac{3}{8} f_e \quad (84)$$

because the inhomogeneity factor of the Earth is given by

$$\Lambda = \frac{C}{M_{\oplus} (a^2 + b^2)} \approx \frac{C}{2M_{\oplus} R_e^2} \approx \frac{1}{6} \quad (85)$$

The most important determinations from gravity measurements since 1915 have been compiled in a table on the following page.

All these gravity measurements are still based on the standard gravity value of Potsdam ( $\varphi = 52^\circ 22.'86$ ;  $\lambda = +13^\circ 4.'06$ ;  $b = 87 \text{ m}$ )

$$g = 981.2740 \text{ gal}$$

obtained by F. Kuhnen and Ph. Furtwangler (Ref. 43). It is necessary to revise the Potsdam system. For the correction of the Potsdam value, the following data are given (Refs. 44 and 45):

P. R. Heyl and G. S. Cook (Wash. D.C.)	: - 20	milligal	}
Bullard (Teddington, G. Brit.)	: - 15	"	
J. S. Clarke (Teddington, Gt. Brit.)	: - 13	"	
Ivanoff (Leningrad, U.S.S.R.)	: - 4	"	
P. R. Heyl	: - 15	"	}
Bullard and Browne	: - 16	"	
Morelli (1954)	: - 16	"	
H. Jeffreys	: - 13.4	"	
Wollard	: - 14 to - 18	"	
A. Berroth	: - 12.5	"	
Morelli (1959)	: - 12.9	"	

Author	Year	Ref.	$g^\circ$ (gal)	$10^6 \beta$	$10^6 \gamma$	$10^6 \delta$	$\lambda_0$	$\Delta R_e$ (m)	$1/f_1$	$1/f_2$	$1/f_m$	$1/f_e$
F. R. Helmert	1915	32	978.052 $\pm 3$	5285 $\pm 5$	18 $\pm$ 3	- 17 $\pm$ 4	230 $\pm$ 51		296.7 $\pm 0.4$	28 000		
A. Berroth	1916	33	978.046	5296		11.6 $\pm$ 4	- 10	150 $\pm$ 58	296.7	297.8 $\pm 0.7$	42 000	
W. Heiskanen	1924	34	978.048 $\pm 3$	5293 $\pm 6$	- 7	---	---	---	---	297.4 $\pm 0.5$	---	
"	1924	34	978.052 $\pm 3$	5285 $\pm 6$	- 7	27 $\pm$ 3	+ 18 $\pm$ 5	345 $\pm$ 38 $\pm 0.6$	294.3 $\pm 0.6$	299.0 $\pm 0.6$	296.7 $\pm 0.5$	18 700
"	1928	35	978.049 $\pm 1$	5289		---	---	---	---	---	297.06	---
"	1928	35	978.049 $\pm 1$	5293		19 $\pm$ 3	0 $\pm$ 5	242	295.7	299.0	297.3	26 703
"	1938	36	978.0451	5302.7		---	---	---	---	---	298.25 $\pm 0.3$	---
N. F. Shuravlev	1938	36	978.0524	5297.0		27.6	- 25	352 $\pm$ 30 $\pm 0.4$	295.3 $\pm 0.4$	300.2 $\pm 0.4$	297.8 $\pm 0.4$	18 314
E. Niskanen	1940	37	978.0484	5303		---	---	---	---	---	298.3	---
H. Jeffreys	1945	38	978.0468	5297.8	5.9	23.0	- 3.9	293	295.7 $\pm 0.2$	299.8 $\pm 0.2$	297.8 $\pm 0.2$	21 550
H. Schutte	1948	39	978.0513	5285.9		---	---	---	---	296.85 $\pm 0.66$	296.3	---
U. A. Uotila	1957	40	978.0520 $\pm 33$	5282.7 $\pm 6.0$		---	---	---	---	297.2	297.4	47 600
Heiskanen & Uotila	1957	41	978.0516	5291.0		10.6	- 6	---	---	297.4	---	
Mean Value			978.049 <sub>s</sub>	5292.4						297.3		

Taking for  $g_e$  the latest determination of Heiskanen and Uotila (Ref. 42) which is nearly in agreement with the average value of all determinations, and using the correction due to H. Jeffreys (Ref. 39) the following value is obtained

$$g_e = 978.0496 - 0.0134 = 978.0362 \text{ gal} = 9.780362 \text{ m/sec}^2 \quad (86)$$

## 9. THE DYNAMIC OBLATENESS AND THE CONSTANTS OF PRECESSION AND NUTATION

The dynamic flattening  $H = (C - A)/C$  is connected with Newcomb's constant of precession,  $P$ , by the relation

$$\frac{P}{H} = 530977.''04 + \frac{94419319''}{1/\kappa + 1} = 1678932.''29 \text{ for } 1/\kappa = 81.250 \quad (87)$$

while the constant of nutation,  $N$ , is given by

$$\frac{N}{H} = \frac{252871''}{1/\kappa + 1} \cos \epsilon = \frac{231982''}{1/\kappa + 1} = 2820.''45 \text{ for } 1/\kappa = 81.250 \quad (88)$$

where  $\cos \epsilon = 0.9173917$  (for 1900.0) has been used for the cosine of the obliquity,  $\epsilon$ , of the ecliptic. The constants in the equations are obtained from Brown's theory of the motion of the Moon and are well known. Both equations yield

$$\frac{P}{N} = 2.288872(1/\kappa + 1) + 407.01140 = 595.271 \text{ for } 1/\kappa = 81.250$$

while observed modern values of  $P$  and  $N$  lead to

$$\frac{P}{N} = \frac{5493.''62}{9.''208} = 596.614 \text{ and thus } 1/\kappa = \frac{P/N}{2.288872} - 178.8218 = 81.84$$

This value for  $1/\kappa$  is by far too large. H. Jeffreys (Ref. 39) has shown that in the equation for the constant of nutation,  $N$ , another constant  $H'$  for the dynamic flattening must be used due to the deviation of the Earth's interior from the isostatic equilibrium ( $H' < H$ ). Therefore  $H'$  can be determined only from  $P$  and  $1/\kappa$ . With  $p_0 = p_e + p_\theta$ , the lunisolar precession,  $p_e = \frac{3}{2} \frac{v_\oplus^2}{c^2} n_\oplus$ , the geodetic precession (a relativistic term due to W. deSitter),  $p$ , the general precession in longitude, and  $\lambda$ , the planetary precession in right ascension, Newcomb's precessional constant is

$$P = \frac{p_0}{\cos \epsilon} = \frac{p + p_\theta}{\cos \epsilon} + \lambda \quad (89)$$

Values for 1900.0 derived from observations are (for a tropical century)

$P = 5490.''66$	$p = 5025.''641$	$\lambda = 12.''473$	$p_\theta = 0$	(Newcomb & Andoyer, Ref. 46)
$5493.156 \pm 0.175$	$5026.000$	$12.493$	$1.915$	(deSitter & Brouwer, Ref. 10)
$5493.847$	$5026.''650$	$12.469$	$1.921$	(Clemence, Ref. 7)

According to newer investigations, Newcomb's value of the general precession in longitude

must be corrected by  $\Delta p = + 0.^{\circ}75$  (H. R. Morgan, Ref. 47),  $\Delta p = + 0.^{\circ}71$  (J. H. Oort, Ref. 48),  $\Delta p = + 0.^{\circ}86$  (Dirk Brouwer, Ref. 48),  $\Delta p = + 0.^{\circ}84$  (Pulkovo Obs., Ref. 48). The average value for the correction may be  $\Delta p = + 0.^{\circ}80$ . All these investigators take  $\Delta\lambda = 0$ . In another paper, J. H. Oort (1943, Ref. 49) takes  $\Delta\lambda = + 0.^{\circ}02$ , a correction also used by deSitter. The correction for  $P$  is therefore

$$\Delta P = \frac{\Delta p + \Delta p_{\text{A}}}{\cos \epsilon} + \Delta\lambda = \frac{0.^{\circ}80 + 1.92}{0.9173917} + 0.00 = 2.^{\circ}96 \quad (90)$$

The value

$$P = 5490.^{\circ}66 + 2.^{\circ}96 = 5493.^{\circ}62 \quad (91)$$

will be accepted here. The dynamic flattening is now

$$H \equiv \frac{C - A}{C} = \frac{5493.^{\circ}62}{1678932.^{\circ}29} = 0.003272091 = \frac{1}{305.615 \pm 0.05} \quad (92)$$

and thus

$$q \equiv \frac{3}{2} \frac{C}{M_{\oplus} R_{\oplus}^2} = \frac{J}{H} = 0.49616_0 \pm 0.00017 \quad (93)$$

and

$$\frac{C}{M_{\oplus} R_{\oplus}^2} = \frac{J_2}{H} = 0.33077_3 \pm 0.00011 \quad (94)$$

The quantity  $q$  may be calculated in another way. Clairaut's theory for the Earth in hydrostatic equilibrium has been developed to the second order by Radau (Ref. 50), Callandreau (Ref. 51), and Darwin (Ref. 28). deSitter (Ref. 31) gives

$$q \equiv \frac{3}{2} \frac{C}{M R_{\oplus}^2} = 1 - \frac{1}{3} \tilde{\omega}_1 - \frac{2}{5} \left(1 - \frac{2}{3} f\right) \frac{\sqrt{1 + \eta_1}}{1 + \lambda_1} \quad (95)$$

where

$$\eta_1 = \frac{\frac{5}{2} \tilde{\omega}_1 + \frac{10}{21} \tilde{\omega}_1^2 + \frac{4}{7} f^2 - \frac{6}{7} f \tilde{\omega}_1}{f - \frac{5}{42} f^2 + \frac{4}{7} \kappa} - 2 \quad (96)$$

and  $1 + \lambda_1$  is an average value of Radau's function,  $f(\eta)$ , depending on the internal density distribution of the Earth. The most reliable value,  $1 + \lambda_1 = 1.00016$ , was derived by Bullard (Ref. 30). With the above data for  $f$  and  $\tilde{\omega}_1$  the above-mentioned equations give, for  $\kappa = 0$ ,

$$\eta_1 = 0.57440 ; q = 0.49815$$

$$\frac{C}{M R_{\oplus}^2} = \frac{2}{3} q = 0.33210$$

These data are not compatible with the previously derived data (eqs. 93 and 94), showing that the hypothesis of hydrostatic equilibrium is not fulfilled for the Earth.

## 10. DENSITY DISTRIBUTION WITHIN THE MOON

It is very difficult to derive a consistent system of lunar constants. Most reports on this subject are based on the work of Sir Harold Jeffreys. However, not even this source is free of inconsistencies. The reason is that many lunar constants are coupled with each other by relations. Therefore a systematic investigation of these relations will be necessary.

It is assumed that the density  $\rho$  within the Moon is constant over concentric ellipsoidal shells

$$\frac{\xi^2}{a^2} + \frac{\eta^2}{b^2} + \frac{\zeta^2}{c^2} = \mu^2 \quad (97)$$

where  $\mu$  varies from 0 at the center to 1 at the surface, and where

$$\begin{aligned}\xi/a &= \mu \cos \phi \cos \theta \\ \eta/b &= \mu \cos \phi \sin \theta \\ \zeta/c &= \mu \sin \phi\end{aligned} \quad (98)$$

are the relative coordinates of the mass element

$$dm = \rho(\mu) d\xi d\eta d\zeta = \rho(\mu) abc \mu^2 \cos \phi d\mu d\phi d\theta \quad (99)$$

The angle  $\phi$  is the lunicentric latitude and  $\theta$  the longitude.  $a$  is the longest semi-axis of the surface ellipsoid pointing toward the Earth,  $b$  the smallest semi-axis in the lunar equator, and  $c$  the rotational or polar semi-axis.

Using equations (98) and (99) after observing that

$$\int_0^{2\pi} \sin^2 \theta d\theta = \int_0^{2\pi} \cos^2 \theta d\theta = \pi ; \int_{-\pi/2}^{\pi/2} \cos^3 \phi d\phi = \frac{4}{3} ; \int_{-\pi/2}^{\pi/2} \sin^2 \phi \cos \phi d\phi = \frac{2}{3}$$

the moments of inertia around the  $a$ ,  $b$ ,  $c$  axes, respectively, become

$$\begin{aligned}A &= \int_0^M (\eta^2 + \zeta^2) dm = \lambda M (b^2 + c^2) \\ B &= \int_0^M (\zeta^2 + \xi^2) dm = \lambda M (c^2 + a^2) \\ C &= \int_0^M (\xi^2 + \eta^2) dm = \lambda M (a^2 + b^2)\end{aligned} \quad (100)$$

where the integrations are taken from 0 to  $2\pi$  with respect to  $\theta$ , from  $-\pi/2$  to  $\pi/2$  with respect to  $\phi$  and from 0 to 1 with respect to  $\mu$ . In the last equation  $M = 4/3 \pi a b c \rho_m$  is the total mass ( $\rho_m$  is the mean density) and the inhomogeneity factor,  $\lambda$ , is given by

$$\lambda = \frac{\int_0^1 \rho \mu^4 d\mu}{\rho_m} = \frac{\int_0^1 \rho \mu^4 d\mu}{3 \int_0^1 \rho \mu^2 d\mu} \quad (101)$$

Because  $a > b > c$ , there is  $A < B < C$ . A constant density model gives  $\lambda = 1/5 = 0.2$ . For Jeffreys compressional model with constant bulk modulus (Ref. 52) Roche's density law [with  $\rho_m = 3.342 \text{ g/cm}^3$  (mean density),  $\rho_0 = 3.290 \text{ g/cm}^3$  (surface density) and  $\rho_c = 3.420 \text{ g/cm}^3$  (central density)] becomes

$$\rho = \rho_c - (\rho_c - \rho_0) \mu^2 = 3.420 - 0.130 \mu^2 [\text{g/cm}^3] \quad (102)$$

The inhomogeneity factor is, therefore,

$$\begin{aligned} \lambda = \frac{1}{5} \frac{\rho_c - \frac{5}{7}(\rho_c - \rho_0)}{\rho_m} &= \frac{1}{5} \frac{\rho_c - \frac{5}{7}(\rho_c - \rho_0)}{\rho_c - \frac{3}{5}(\rho_c - \rho_0)} = \frac{0.9955}{5} = \\ &= 0.1991 \pm 0.0001 \end{aligned} \quad (103)$$

## 11. CONSTANTS OF THE PHYSICAL LIBRATION OF THE MOON

The values of  $f \equiv \frac{C-B}{C-A}$  and the inclination of the Moon's equator to the ecliptic can be

determined from observations of the physical libration of the Moon. Due to the difficulty of observations near the irregular limb of varied illumination the values for  $f$  scatter widely, as can be seen from the following table (Refs. 11 and 13):

Author	Year	$f \equiv \frac{C-B}{C-A}$
F. Hayn	1907	0.75 $\pm$ 0.04
F. Hayn		0.85 $\pm$ 0.07
J. Stratton	1909	0.50 $\pm$ 0.03
I. V. Belkovich	1936	0.84 $\pm$ 0.08
I. V. Belkovich	1949	0.67 $\pm$ 0.03
K. Koziel	1949	0.71 $\pm$ 0.051
K. Koziel	1949	0.60 $\pm$ 0.055
A.A. Nefedjev	1950	0.65 $\pm$ 0.045
A.A. Yakovkin	1950	0.85 $\pm$ 0.03
T. Weimer	1954	0.60
Mean Value		0.70 <sub>2</sub>

Sir Harold Jeffreys used  $f = 0.84$  in his book *The Earth* (Ref. 11). Later he recommended  $f = 0.67$  (Ref. 53) and used  $f = 0.639 \pm 0.014$  in his latest paper (Ref. 54).

The secular motions of the perigee and node of the lunar orbit are also influenced by the Moon's oblateness coefficients ( $L$  and  $K$ ). From Jeffreys equation for the perigee motion follows (Ref. 11):

$$380L - 1192K = 6.420 - 3896J_\oplus \geq 0$$

and thus

$$f = 1 - \frac{K}{L} \geq 1 - \frac{380}{1192} = \frac{812}{1192} = 0.6812$$

The mean value of the table is consistent with this lower limit for  $f$  and therefore

$$f = 0.70 \pm 0.02 \quad (104)$$

will be adopted in this paper.

A new investigation of the libration of the Moon's axis by H. Jeffreys (Ref. 54) leads to

$$\beta \equiv \frac{C - A}{C} = 0.0006279 \pm 0.0000010 \quad (105)$$

taking into account a solar effect not evaluated by Hayn.

The three quantities  $\lambda$ ,  $f$ , and  $\beta$  are sufficient to calculate all other quantities, provided the mass and the mean radius of the Moon are known.

## 12. RELATIONS AND NUMERICAL VALUES FOR THE DIMENSIONLESS MOMENT OF INERTIA PARAMETERS

The same symbols for moment of inertia parameters will be used as they have been introduced mainly by H. Jeffreys (Ref. 11). The numerical values are based on the above-given parameters  $\lambda$ ,  $f$ , and  $\beta$ ; namely

$$\begin{aligned} \lambda &= \frac{A}{M(b^2 + c^2)} = \frac{B}{M(c^2 + a^2)} = \frac{C}{M(a^2 + b^2)} \\ &= \frac{C - B}{M(b^2 - c^2)} = \frac{C - A}{M(a^2 - c^2)} = \frac{C - B}{M(b^2 - c^2)} = 0.1991 \pm 0.0001 \end{aligned} \quad (106)$$

$$f = \frac{\alpha}{\beta} = \frac{C - B}{C - A} = \frac{2J - K}{2J + K} = \frac{J - 1/2 K}{L} = \frac{b^2 - c^2}{a^2 - c^2} = 0.70 \pm 0.02 \quad (107)$$

$$\beta = \frac{C - A}{C} = \frac{J + 1/2 K}{g} = \frac{L}{g} = \frac{a^2 - c^2}{a^2 + b^2} = 0.0006279 \pm 0.0000010 \quad (108)$$

The other parameters can be derived from these as follows:

$$\begin{aligned} 1 - f &= \frac{\gamma}{\beta} = \frac{B - A}{C - A} = \frac{K}{J + 1/2 K} = \frac{K}{L} = \frac{a^2 - b^2}{a^2 - c^2} = \frac{3\lambda - g}{L} = \frac{3\lambda/g - 1}{\beta} = \\ &= 0.30 \pm 0.02 \end{aligned} \quad (109)$$

$$\alpha = \frac{C - B}{C} = f\beta = \frac{J - 1/2 K}{g} = \frac{L - K}{g} = \frac{b^2 - c^2}{a^2 + b^2} = 0.0004395 \pm 0.0000133$$

(110)

$$\gamma = \frac{B - A}{C} = \beta - \alpha = \frac{K}{g} = (1 - f)\beta = \frac{a^2 - b^2}{a^2 + b^2} = 0.0001884 \pm 0.0000129$$

(111)

$$g = \frac{3}{2} \frac{C}{M a^2} = \frac{3\lambda}{1 + \gamma} = \frac{3}{2} \lambda \frac{a^2 + b^2}{a^2} = 0.5972 \pm 0.0003$$

(112)

$$L = \frac{3}{2} \frac{C - A}{M a^2} = J + \frac{1}{2} K = g\beta = \frac{3}{2} \lambda \frac{a^2 - c^2}{a^2} = 0.0003750 \pm 0.0000008$$

(113)

$$J = \frac{3}{2} \frac{C - \frac{A + B}{2}}{M a^2} = L - \frac{1}{2} K = \frac{1 + f}{2} L = \frac{1}{2} \frac{1 + f}{1 - f} K = \frac{3}{2} \lambda \frac{\frac{a^2 + b^2}{2} - c^2}{a^2}$$

$$= g \left( \beta - \frac{1}{2} \gamma \right) = g \frac{a + \beta}{2} = g \left( a + \frac{1}{2} \gamma \right) = 0.0003187 \pm 0.0000044$$

(114)

$$K = \frac{3}{2} \frac{B - A}{M a^2} = (1 - f)L = g\gamma = \frac{3}{2} \lambda \frac{a^2 - b^2}{a^2} = 0.0001125 \pm 0.0000077$$

(115)

The dimensionless moments of inertia and their differences are obtained from the above-mentioned data as:

$$\frac{C - A}{M a^2} = \frac{2}{3} L = \frac{2J + K}{3} = \lambda \frac{a^2 - c^2}{a^2} = 0.0002500 \pm 0.0000005$$

(116)

$$\frac{C - B}{M a^2} = \frac{2}{3} fL = \frac{2J - K}{3} = \lambda \frac{b^2 - c^2}{a^2} = 0.0001750 \pm 0.0000054$$

(117)

$$\frac{B - A}{M a^2} = \frac{2}{3} (1 - f)L = \frac{2}{3} K = \lambda \frac{a^2 - b^2}{a^2} = 0.0000750 \pm 0.0000051$$

(118)

$$\frac{A}{M a^2} = \frac{2}{3} g (1 - \beta) = \frac{2}{3} (g - L) = \lambda \frac{b^2 + c^2}{a^2} = 0.3978_{77} \pm 0.0002$$

(119)

$$\frac{B}{M a^2} = \frac{2}{3} g (1 - \alpha) = \frac{2}{3} (g - fL) = \lambda \frac{c^2 + a^2}{a^2} = 0.3979_{52} \pm 0.0002$$

(120)

$$\frac{C}{M\alpha^2} = \frac{2}{3} g = \frac{2\lambda}{1+\gamma} = \lambda \frac{\alpha^2 + b^2}{\alpha^2} = 0.3981_{27} \pm 0.0002 \quad (121)$$

The ratios of the semi-axes now become

$$\frac{b}{a} = \sqrt{\frac{1-\gamma}{1+\gamma}} = \sqrt{1 - \frac{2\gamma}{1+\gamma}} = 0.9998116 \pm 0.0000129 \quad (122)$$

$$\frac{c}{a} = \sqrt{1 - \frac{2\beta}{1+\gamma}} = \sqrt{1 - \frac{2\gamma}{(1-f)(1+\gamma)}} = 0.9993720 \pm 0.0000010 \quad (123)$$

These data seem, at present, to be the most reasonable. H. Jeffreys' value

$$g \equiv \frac{3}{2} \frac{C}{M\alpha^2} = 0.5956 \pm 0.0010$$

is slightly low, and affords a higher density concentration towards the center as has been assumed by Jeffreys.

### 13. THE FINAL DETERMINATION OF THE DIMENSIONS, MASSES AND MOMENTS OF INERTIA FOR THE EARTH AND THE MOON

Using the obtained value of the gravitational acceleration,  $g_e$ , at the equator (eq. 86) in the relations (eq. 80) and (eq. 82) at the end of Section 7 there follows at once  $R_e = \sqrt{\chi \mu_\oplus / g_e} = 6378169.835$  m and  $\rho_\oplus = 5.516964$  g/cm<sup>3</sup>. The final values, adopted for the Earth, will be taken as

$$R_e + 6378170 (1 \pm 3.2 \times 10^{-6}) \text{ m} = 6378170 \pm 20 \text{ m} \quad (124)$$

and

$$\rho_\oplus = 5.5170 (1 \pm 7.3 \times 10^{-4}) \text{ g/cm}^3 = 5.5170 \pm 0.0040 \text{ g/cm}^3 \quad (125)$$

The volume of the Earth is

$$\begin{aligned} V_\oplus &= \frac{4}{3} \pi R_e^3 (1-f) = 1.083225 \times 10^{27} (1 \pm 1.02 \times 10^{-5}) \text{ cm}^3 \\ &= (1.083225 \pm 0.000011) \times 10^{27} \text{ cm}^3 \end{aligned} \quad (126)$$

while the mass is given by

$$M_\oplus = \frac{\mu_\oplus}{G} = V_\oplus \rho_\oplus = 5.9761 \times 10^{27} (1 \pm 7.2 \times 10^{-4}) \text{ g} = (5.9761 \pm 0.0043) \times 10^{27} \text{ g} \quad (127)$$

and the polar radius now becomes

$$R_p = R_e (1-f) = 6356788 (1 \pm 3.7 \times 10^{-6}) \text{ m} = 6356788 \pm 24 \text{ m} \quad (128)$$

The unit for the moments of inertia is

$$M_{\oplus} R_{\oplus}^2 = 2.4311_4 \times 10^{45} (1 \pm 7.3 \times 10^{-4}) g \text{ cm}^2 = (2.4311_4 \pm 0.0018) \times 10^{45} g \text{ cm}^2 \quad (129)$$

and therefore

$$C - A = J_2 \times M_{\oplus} R_{\oplus}^2 = 2.6313 \times 10^{42} (1 \pm 9.1 \times 10^{-4}) g \text{ cm}^2 = (2.6313 \pm 0.0024) \times 10^{42} g \text{ cm}^2 \quad (130)$$

$$C = \frac{2}{3} q \times M_{\oplus} R_{\oplus}^2 = 8.0415 \times 10^{44} (1 \pm 1.07 \times 10^{-3}) g \text{ cm}^2 = (8.0415 \pm 0.0086) \times 10^{44} g \text{ cm}^2 \quad (131)$$

It is now possible to give corresponding data for the Moon. The mass is given by

$$M_{\oplus} = \kappa M_{\oplus} = 7.3552 \times 10^{25} (1 \pm 1.02 \times 10^{-3}) g = (7.3552 \pm 0.0075) \times 10^{25} g \quad (132)$$

while the mean visible radius is

$$R_{\oplus} = k R_{\oplus} = 1738236 (1 \pm 1.0 \times 10^{-4}) = 1738236 \pm 174 \text{ m} \quad (133)$$

and therefore the semi-axes of the three-axial Moon are

$$a = \frac{R_{\oplus}}{\sigma} = 1738946 \pm 186 \text{ m} \quad (134)$$

$$b = a \left( \frac{b}{a} \right) = 1738618 \pm 209 \text{ m} \quad (135)$$

$$c = a \left( \frac{c}{a} \right) = 1737854 \pm 188 \text{ m} \quad (136)$$

The unit for the Moon's moments of inertia is

$$M_{\oplus} a^2 = 2.2241_6 \times 10^{42} (1 \pm 1.23 \times 10^{-3}) g \text{ cm}^2 = (2.2241_6 \pm 0.0027) \times 10^{42} g \text{ cm}^2 \quad (137)$$

and thus the moments of inertia are

$$A = 0.8849_{42} \times 10^{42} (1 \pm 1.73 \times 10^{-3}) g \text{ cm}^2 = (0.8849_{42} \pm 0.0015_3) \times 10^{42} g \text{ cm}^2 \quad (138)$$

$$B = 0.8851_{09} \times 10^{42} (1 \pm 1.73 \times 10^{-3}) g \text{ cm}^2 = (0.8851_{09} \pm 0.0015_3) \times 10^{42} g \text{ cm}^2 \quad (139)$$

$$C = 0.8854_{98} \times 10^{42} (1 \pm 1.73 \times 10^{-3}) g \text{ cm}^2 = (0.8854_{98} \pm 0.0015_3) \times 10^{42} g \text{ cm}^2 \quad (140)$$

$$C - \frac{A + B}{2} = (0.000473 \pm 0.000007) \times 10^{42} g \text{ cm}^2 \quad (141)$$

$$B - A = (0.000167 \pm 0.000012) \times 10^{42} g \text{ cm}^2 \quad (142)$$

The oblateness coefficients of the potential function of the Moon are

$$J_2 = \frac{C - (A + B)/2}{M_{\oplus} a^2} = \frac{2}{3} J = 0.0002125 \pm 0.0000029 \quad (143)$$

$$J_2^{(2)} = \frac{B - A}{4 M_e a^2} = \frac{1}{6} K = 0.0000188 \pm 0.0000013 \quad (144)$$

The derived value for the equatorial radius of the Earth (eq. 124) is in good agreement with the following values:

Author	Year	Ref.	$R_e$
W. M. Kaula	1961	55	$6378163 \pm 21$ m
V. C. Clarke, Jr.	1962	56	$6378165 \pm 25$
I. Fischer	1962	57	$6378166$
Present Report	1962		$6378170 \pm 20$

I. Fisher's value for  $1/\kappa = M_\oplus/M_e = 81.268$  is also in good agreement with the value in this report. The presented system of constants is not only a consistent one, but the most serious discrepancy has been removed in determining the gravitational parameter  $\mu_\oplus$  from terrestrial data and, on the other hand, from the lunar mean motion in combination with radar measurements of the Moon's distance.

Finally, the present data for the Moon's moments of inertia are compared with the values of other authors in the following table:

Author	Ref.	$A$ $10^{35} \text{kgm}^2$	$B$ $10^{35} \text{kgm}^2$	$C$ $10^{35} \text{kgm}^2$	$C - (A+B)/2$ $10^{35} \text{kgm}^2$	$B - A$ $10^{35} \text{kgm}^2$
B. E. Kalensher	58	0.87976	0.87985	0.88032	0.00051	0.00009
Makemson, Baker, Westrom	13	0.88837	0.88856	0.88893	0.00047	0.00019
V. C. Clarke, Jr.	56	0.88746	0.88764	0.88801	0.00046	0.00018
Present Report	---	0.88494	0.88511	0.88550	0.00047	0.00017

The values of V. C. Clarke, Jr. are used for the Ranger Program.

#### 14. THE EARTH ELLIPSOID

The equation of the rotational ellipsoid or spheroid is

$$\frac{x^2 + y^2}{R_e^2} + \frac{z^2}{R_p^2} = 1 \quad (145)$$

where

$$x = R \cos \phi \cos \lambda = R_e \cos \psi \cos \lambda = \rho_n \cos \varphi \cos \lambda \quad (146)$$

$$y = R \cos \phi \sin \lambda = R_e \cos \psi \sin \lambda = \rho_n \cos \varphi \sin \lambda \quad (147)$$

$$z = R \sin \phi = R_p \sin \psi = \rho_n (1 - e^2) \sin \varphi \quad (148)$$

and  $R_e$  is the equatorial Earth radius,  $R_p$  the polar Earth radius,  $R$  the local Earth radius,  $\rho_n$  the normal radius of curvature,  $\lambda$  the geographic longitude (positive eastward of Greenwich),  $\phi$  the geocentric latitude,  $\psi$  the reduced latitude,  $\varphi$  the geodetic or geographic ( $\approx$  astronomical) latitude, and  $e$  the first eccentricity of the meridian ellipse. Introducing the second eccentricity,  $\epsilon$ , and the flattening (oblateness, ellipticity),  $f$ , the following relations hold

$$f = \frac{R_e - R_p}{R_e} = 1 - \sqrt{1 - e^2} = 1 - \frac{1}{\sqrt{1 + \epsilon^2}} \approx \frac{1}{2} e^2 + \frac{1}{8} e^4 + \dots + \frac{1 \cdot 1 \cdot 3 \cdot 5 \cdots (2\kappa - 3)}{2 \cdot 4 \cdot 6 \cdot 8 \cdots (2\kappa)} e^{2\kappa} + \dots \quad (149)$$

$$\epsilon^2 = \frac{R_e^2 - R_p^2}{R_e^2} = \frac{\epsilon^2}{1 + \epsilon^2} = f(2 - f) = 2f - f^2 \quad (150)$$

$$\epsilon^2 = \frac{R_e^2 - R_p^2}{R_e^2} = \frac{\epsilon^2}{1 - e^2} = \frac{f(2 - f)}{(1 - f)^2} = 2f + 3f^2 + \dots + (\kappa + 1)f^\kappa + \dots \quad (151)$$

thus

$$\frac{R_p}{R_e} = 1 - f = \sqrt{1 - e^2} = \frac{1}{\sqrt{1 + \epsilon^2}} \quad (152)$$

The different latitude angles are related by

$$\tan \phi = \sqrt{1 - e^2} \tan \psi = (1 - e^2) \tan \varphi \quad (153)$$

$$\tan \phi = (1 - f) \tan \psi = (1 - f)^2 \tan \varphi \quad (154)$$

By differentiation the relation

$$\frac{R^2}{R_p} d\phi = R_e d\psi = \rho d\varphi \quad (155)$$

follows.

Thus the line element is

$$ds^2 = dx^2 + dy^2 + dz^2 = dr^2 + r^2 d\phi^2 + r^2 \cos^2 \phi d\lambda^2 = R_e^2 (1 - e^2 \cos^2 \psi) d\psi^2 + R_e^2 \cos^2 \psi d\lambda^2 \quad (156)$$

The parameter  $\rho$  is the mean radius of curvature, and is correlated with the normal radius of curvature

$$\begin{aligned} \rho_n &= \frac{R_e}{(1 - e^2 \sin^2 \varphi)^{1/2}} = \frac{R_e^2 / R_p}{(1 + \epsilon^2 \cos^2 \varphi)^{1/2}} = R \frac{\cos \phi}{\cos \varphi} = \\ &= R_e \left[ 1 + \frac{1}{2} e^2 \sin^2 \varphi + \frac{3}{8} e^4 \sin^4 \varphi + \frac{5}{16} e^6 \sin^6 \varphi + \dots \right] \end{aligned} \quad (157)$$

and to the meridional radius of curvature

$$\rho_m = \frac{R_e (1 - e^2)}{(1 - e^2 \sin^2 \varphi)^{3/2}} = \frac{R_p^2 / R_e}{(1 - e^2 \sin^2 \varphi)^{3/2}} = \frac{R_e^2 / R_p}{(1 + e^2 \cos^2 \varphi)^{3/2}}$$

$$= R_e (1 - e^2) \left[ 1 + \frac{3}{2} e^2 \sin^2 \varphi + \frac{15}{8} e^4 \sin^4 \varphi + \frac{35}{16} e^6 \sin^6 \varphi + \dots \right] \quad (158)$$

by the relation

$$\rho = \sqrt{\rho_m \rho_n} = \frac{R_p}{1 - e^2 \sin^2 \varphi} = \frac{R_e^2 / R_p}{1 + e^2 \cos^2 \varphi} = R_p [1 + e^2 \sin^2 \varphi + e^4 \sin^4 \varphi + e^6 \sin^6 \varphi + \dots] \quad (159)$$

The radius of a parallel of latitude is  $\rho_n \cos \varphi = R_e \cos \psi = R \cos \phi$ . Because  $f$  or  $e^2$  are small quantities the latitudes  $\phi$ ,  $\psi$ , and  $\varphi$  will not differ very much from each other. Therefore it is very useful to have rapidly converging series developments available for the differences  $\phi - \varphi$  and  $\psi - \varphi$ . With

$$m = \frac{R_e^2 - R_p^2}{R_e^2 + R_p^2} = \frac{1 - (1 - e^2)}{1 + (1 - e^2)} = \frac{e^2}{2 - e^2} = f + \frac{1}{2} f^2 - \frac{1}{4} f^4 + \dots \quad (160)$$

there is

$$\phi = \varphi - m \sin 2 \varphi + \frac{m^2}{2} \sin 4 \varphi - \frac{m^3}{3} \sin 6 \varphi \pm \dots \quad (161)$$

$$\varphi = \phi + m \sin 2 \phi + \frac{m^2}{2} \sin 4 \phi + \frac{m^3}{3} \sin 6 \phi + \dots \quad (162)$$

and with

$$n = \frac{R_e - R_p}{R_e + R_p} = \frac{1 - (1 - f)}{1 + (1 - f)} = \frac{f}{2 - f} = \frac{1}{2} f + \frac{1}{4} f^2 + \dots \quad (163)$$

there is

$$\psi = \varphi - n \sin 2 \varphi + \frac{n^2}{2} \sin 4 \varphi - \frac{n^3}{3} \sin 6 \varphi \pm \dots \quad (164)$$

$$\varphi = \psi + n \sin 2 \psi + \frac{n^2}{2} \sin 4 \psi + \frac{n^3}{3} \sin 6 \psi + \dots \quad (165)$$

An accurate formula for the difference  $\varphi - \phi$  is given by

$$\tan(\varphi - \phi) = \frac{e^2 \tan \varphi}{1 + (1 - e^2) \tan^2 \varphi} = \frac{e^2 \sin \varphi \cos \varphi}{1 - e^2 \sin^2 \varphi} \quad (166)$$

The local Earth radius (radius vector) can be accurately calculated from the relations

$$R = \sqrt{x^2 + y^2 + z^2} = \frac{R_p}{\sqrt{1 - e^2 \cos^2 \phi}} = R_e \sqrt{1 - e^2 \sin^2 \psi} = R_e \sqrt{\frac{1 - e^2 (2 - e^2) \sin \varphi}{1 - e^2 \sin^2 \varphi}}$$

$$= R_e \sqrt{\frac{1 - 4m(1+m)^{-2} \sin^2 \varphi}{1 - 4n(1+n)^{-2} \sin^2 \varphi}} = \frac{1+n}{1+m} R_e \sqrt{\frac{1 + m^2 + 2m \cos 2\varphi}{1 + n^2 + 2n \cos 2\varphi}} \quad (167)$$

By taking the logarithm of the last relation there follows

$$\ln \frac{R}{R_e} = \ln \frac{1+n}{1+m} + \frac{1}{2} \ln (1+m^2 + 2m \cos 2\varphi) - \frac{1}{2} \ln (1+n^2 + 2n \cos 2\varphi)$$

or, using a known series development,

$$\log \frac{R}{R_e} = \log \frac{1+n}{1+m} + M \left[ (m-n) \cos 2\varphi - \frac{m^2-n^2}{2} \cos 4\varphi + \frac{m^3-n^3}{3} \cos 6\varphi + \dots \right] \quad (168)$$

where  $M$  is the module ( $M = 0.4342944819$ ). This series is due to Encke.

Conventional power series for  $R/R_e$  can be obtained as follows:

$$\frac{R}{R_e} = (1+\epsilon^2 \sin^2 \phi)^{-1/2} = 1 - \frac{1}{2} \epsilon^2 \sin^2 \phi + \frac{3}{8} \epsilon^4 \sin^4 \phi - \frac{5}{16} \epsilon^6 \sin^6 \phi + \dots \quad (169)$$

The above-mentioned power series for the local Earth radius can also be written

$$\frac{R}{R_e} = 1 - \left( \frac{1}{2} \epsilon^2 - \frac{3}{8} \epsilon^4 + \frac{5}{16} \epsilon^6 \right) \sin^2 \phi - \left( \frac{3}{32} \epsilon^4 - \frac{5}{64} \epsilon^6 \right) \sin^2 2\phi + \frac{5}{64} \epsilon^6 \sin^2 \phi \sin^2 2\phi + \dots \quad (170)$$

or

$$\frac{R}{R_e} = 1 - f \sin^2 \phi - \left( \frac{3}{8} f^2 + \frac{1}{2} f^3 \right) \sin^2 2\phi + \frac{5}{8} f^3 \sin^2 \phi \sin^2 2\phi + \dots \quad (171)$$

using eq. (151).

In order to obtain power series for the local Earth radius as a function of the geodetic latitude,  $\varphi$ , it is useful to set

$$k = \frac{4m}{(1+m)^2} = e^2 (2 - e^2) ; \quad k' = \frac{4n}{(1+n)^2} = f (2 - f) = e^2 ; \quad k/k' = 2 - e^2 \quad (172)$$

the non-dimensional local Earth radius is now

$$\frac{R}{R_e} = 1 - \frac{1}{2} (k - k') \sin^2 \varphi - \frac{1}{8} (k^2 + 2kk' - 3k'^2) \sin^4 \varphi - \frac{1}{16} (k^3 + k^2 k' + 3kk'^2 - 5k'^3) \sin^6 \varphi - \dots \quad (173)$$

or

$$\begin{aligned} \frac{R}{R_e} = 1 &- \frac{1}{2} e^2 [(2 - e^2) - 1] \sin^2 \varphi - \frac{1}{8} e^4 [(2 - e^2)^2 + 2(2 - e^2) - 3] \sin^4 \varphi \\ &- \frac{1}{16} e^6 [(2 - e^2)^3 + (2 - e^2)^2 + 3(2 - e^2) - 5] \sin^6 \varphi - \dots \end{aligned} \quad (174)$$

Because

$$e^2 = 2/f - f^2 ; \quad 2 - e^2 = 2 - 2f + f^2$$

there is, also,

$$\begin{aligned} \frac{R}{R_e} &= 1 - \left( f - \frac{5}{2} f^2 + 2f^3 \right) \sin^2 \varphi - \left( \frac{5}{2} f^2 - \frac{17}{2} f^3 \right) \sin^4 \varphi - \frac{13}{2} f^3 \sin^6 \varphi - \dots \\ &= 1 - f \sin^2 \varphi + \left( \frac{5}{8} f^2 - \frac{1}{2} f^3 \right) \sin^2 2\varphi + \frac{13}{8} f^3 \sin^2 \varphi \sin^2 2\varphi + \dots \\ &= 1 - \frac{1}{2} f + \frac{5}{16} f^2 + \frac{5}{32} f^3 + \left( \frac{1}{2} f - \frac{13}{64} f^3 \right) \cos 2\varphi - \left( \frac{5}{16} f^2 + \frac{5}{32} f^3 \right) \cos 4\varphi + \frac{13}{64} f^3 \cos 6\varphi + \dots \end{aligned} \quad (175)$$

It is also important to know the arc corresponding to  $1^\circ$  in longitude, namely

$$\begin{aligned} \nu &= \frac{\pi}{180} \rho_n \cos \varphi = \frac{\pi}{180} \frac{R_e \cos \varphi}{(1 - e^2 \sin^2 \varphi)^{1/2}} \\ &= \frac{\pi}{180} R_e \cos \varphi \left( 1 + \frac{1}{2} e^2 \sin^2 \varphi + \frac{3}{8} e^4 \sin^4 \varphi + \dots \right) \\ &= \frac{\pi}{180} R_e \left( 1 + \frac{1}{8} e^2 + \frac{3}{64} e^4 \right) \cos \varphi - \left( \frac{e^2}{8} + \frac{9}{128} e^4 \right) \cos 3\varphi + \frac{3}{128} e^4 \cos 5\varphi + \dots \end{aligned} \quad (176)$$

and the arc corresponding to  $1^\circ$  in latitude, namely

$$\begin{aligned} \mu &= \frac{\pi}{180} \rho_m = \frac{\pi}{180} \frac{R_e (1 - e^2)}{(1 - e^2 \sin^2 \varphi)^{3/2}} = \frac{\pi}{180} R_e (1 - e^2) \left[ 1 + \frac{3}{2} e^2 \sin^2 \varphi + \frac{15}{8} e^4 \sin^4 \varphi + \dots \right] \\ &= \frac{\pi}{180} \frac{R_p^2}{R_e} \left[ 1 + \frac{3}{4} e^2 + \frac{45}{64} e^4 - \left( \frac{3}{4} e^2 + \frac{15}{16} e^4 \right) \cos 2\varphi + \frac{15}{64} e^4 \cos 4\varphi + \dots \right] \end{aligned} \quad (177)$$

The length of a meridian quadrant is given by

$$\begin{aligned} Q_m &= \frac{\pi}{2} \overline{\rho}_m = \int_0^{\pi/2} \frac{R_e (1 - e^2)}{(1 - e^2 \sin^2 \varphi)^{3/2}} d\varphi = R_e (1 - e^2) \int_0^1 \frac{dz}{(1 - e^2 z^2) \sqrt{(1 - z^2)(1 - e^2 z^2)}} \\ &= R_e (1 - e^2) \prod \left( \frac{\pi}{2}, -e^2, e \right) = R_e (1 - e^2) \int_0^{\pi/2} \left[ 1 + \frac{3}{2} e^2 \sin^2 \varphi + \frac{15}{8} e^4 \sin^4 \varphi + \frac{35}{16} e^6 \sin^6 \varphi + \dots \right] d\varphi \\ &= \frac{\pi}{2} R_e (1 - e^2) \left[ 1 + \frac{3}{4} e^2 + \frac{45}{64} e^4 + \frac{175}{256} e^6 + \dots \right] \\ &= \frac{\pi}{2} R_e \left( 1 - \frac{1}{4} e^2 - \frac{3}{64} e^4 - \frac{5}{256} e^6 - \dots \right) \\ &= \frac{\pi}{2} R_e \left( 1 - \frac{1}{2} f + \frac{1}{16} f^2 + \frac{1}{32} f^3 + \dots \right) \end{aligned} \quad (178)$$

The volume of the Earth spheroid is simply given by

$$V = \frac{4}{3} \pi R_e^2 R_p = \frac{4}{3} \pi R_e^3 (1 - f) \quad (179)$$

while the surface area follows from

$$\begin{aligned} S &= 4\pi R_e R_p \int_0^1 \sqrt{1 + \epsilon^2 z^2} dz = 2\pi R_e R_p \left[ \sqrt{1 + \epsilon^2} + \frac{1}{\epsilon} \sinh^{-1} \epsilon \right] \\ &= 2\pi R_e^2 \left[ 1 + \frac{\sinh^{-1} \epsilon}{\epsilon \sqrt{1 + \epsilon^2}} \right] = 2\pi R_e^2 \left[ 1 + \frac{\ln(\epsilon + \sqrt{1 + \epsilon^2})}{\epsilon \sqrt{1 + \epsilon^2}} \right] \\ &= 2\pi R_e^2 \left[ 1 + \left( 1 - \frac{1}{6} \epsilon^2 + \frac{3}{40} \epsilon^4 - \frac{5}{112} \epsilon^6 \pm \dots \right) \left( 1 - \frac{1}{2} \epsilon^2 + \frac{3}{8} \epsilon^4 - \frac{5}{16} \epsilon^6 \pm \dots \right) \right] \\ &= 4\pi R_e^2 \left( 1 - \frac{1}{3} \epsilon^2 + \frac{4}{15} \epsilon^4 - \frac{8}{35} \epsilon^6 - \dots \right) = 4\pi R_e^2 \left( 1 - \frac{2}{3} f + \frac{1}{15} f^2 + \frac{4}{105} f^3 + \dots \right) \end{aligned} \quad (180)$$

These formulas will now be used to derive the Earth's dimensions and other Earth parameters.

Taking  $R_e = 6378170 \pm 20$  m and  $1/f = 298.30 \pm 0.05$  yields

$$e^2 = 0.006693422, \quad e = 0.081813334$$

$$\epsilon^2 = 0.006738525, \quad \epsilon = 0.082088522$$

$$m = 0.003357949 = 692.''627$$

$$n = 0.001678979 = 346.''314$$

The mean radius is

$$\bar{R} = \frac{2R_e + R_p}{3} = 6371043 \pm 21 \text{ m} \quad (181)$$

The radius for the geodetic latitude  $\varphi = \sin^{-1} \sqrt{1/3} = 35^\circ 15' 51.''8$  ( $\phi = 35^\circ 4' 59.''5$ ) is

$$R_1 = R_e \sqrt{1 - \frac{e^2 (1 - e^2)}{3 - e^2}} = R_e \left[ 1 - \frac{1}{3} f + \frac{5}{9} f^2 + \frac{1}{3} f^3 + \dots \right] = 6371083 \pm 21 \text{ m} \quad (182)$$

The radius for the geocentric latitude  $\phi = \sin^{-1} \sqrt{1/3} = 35^\circ 15' 51.''8$  ( $\varphi = 35^\circ 26' 45.''5$ ) is

$$R_2 = R_e \sqrt{1 - \frac{e^2}{3 - 2e^2}} = R_e \left[ 1 - \frac{1}{3} f - \frac{1}{3} f^2 - \frac{7}{27} f^3 - \dots \right] = 6371019 \pm 21 \text{ m} \quad (183)$$

The radius for a sphere of equal area is

$$R_s = (S/4\pi)^{1/2} = R_e \left( 1 - \frac{1}{3} f^2 - \frac{1}{45} f^3 + \frac{11}{945} f^4 + \dots \right) = 6371041 \pm 21 \text{ m} \quad (184)$$

while the radius for a sphere of equal volume is

$$R_V = (3V/4\pi)^{1/3} = R_e (1 - f)^{1/3} = R_e \left( 1 - \frac{1}{3} f - \frac{1}{9} f^2 - \frac{5}{81} f^3 - \dots \right) = 6371035 \pm 21 \text{ m}$$

The surface area is

$$S = 4\pi R_e^2 \left( 1 - \frac{2}{3} f + \frac{1}{15} f^2 + \frac{4}{105} f^3 + \dots \right) = (5.100711 \pm 0.000034) \times 10^{14} \text{ m}^2 \quad (186)$$

while the volume is given by

$$V = \frac{4}{3}\pi R_e^3 (1 - f) = (1.083225 \pm 0.000011) \times 10^{21} \text{ m}^3 \quad (187)$$

The radius of curvature at the pole is

$$\rho_p = R_e^2 / R_p = R_e / (1 - f) = 6399624 \pm 24 \text{ m} \quad (188)$$

while the meridional radius of curvature at the equator is given by

$$\rho_{m_e} = R_p^2 / R_e = R_e (1 - f)^2 = R_e (1 - e^2) = 6335478 \pm 27 \text{ m} \quad (189)$$

The length of an equatorial quadrant is

$$Q_e = \frac{\pi}{2} R_e = 10018806 \pm 31 \text{ m} \quad (190)$$

while the length of a meridional quadrant is

$$Q_m = \frac{\pi}{2} R_e \left( 1 - \frac{1}{2} f + \frac{1}{16} f^2 + \frac{1}{32} f^3 + \dots \right) = 10002020 \pm 34 \text{ m} \quad (191)$$

Therefore the arc corresponding to  $1^\circ$  in longitude is at the equator

$$\nu_e = Q_e / 90 = \frac{\pi}{180} R_e = 111320.07 \pm 0.35 \text{ m} \quad (192)$$

while the arc corresponding to  $1^\circ$  in latitude is at the equator

$$\mu_e = \frac{\pi}{180} \rho_{m_e} = \frac{\pi}{180} R_e (1 - f)^2 = 110574.95 \pm 0.47 \text{ m} \quad (193)$$

at the pole

$$\mu_p = \frac{\pi}{180} \rho_p = \frac{\pi}{180} R_e / (1 - f) = 111694.51 \pm 0.41 \text{ m} \quad (194)$$

and in the average

$$\bar{\mu} = Q_m / 90 = \frac{\pi}{180} R_e \left( 1 - \frac{1}{2} f + \frac{1}{16} f^2 + \frac{1}{32} f^3 + \dots \right) = 111133.56 \pm 0.38 \text{ m} \quad (195)$$

Finally, a few series developments are given for the Earth radius, for the various definitions of latitude, and for the radii of curvature:

$$\begin{aligned}\phi &= \varphi - 692.''627 \sin 2\varphi + 1.''163 \sin 4\varphi \pm \dots \\ \varphi &= \phi + 692.''627 \sin 2\phi + 1.''163 \sin 4\phi + \dots\end{aligned}\quad (196)$$

and

$$\begin{aligned}\psi &= \varphi - 346.''314 \sin 2\varphi + 0.''291 \sin 4\varphi \pm \dots \\ \varphi &= \psi + 346.''314 \sin 2\psi + 0.''291 \sin 4\psi + \dots\end{aligned}\quad (197)$$

furthermore

$$\log \frac{R}{R_e} = 9.99927266 + 0.00072917 \cos 2\varphi - 0.00000184 \cos 4\varphi + \dots \quad (198)$$

and

$$\begin{aligned}\frac{R}{R_e} &= 1 - 0.003369263 \sin^2 \phi + 0.000017028 \sin^4 \phi - 0.000000096 \sin^6 \phi \pm \dots \\ &= 1 - 0.003352330 \sin^2 \phi - 0.000004233 \sin^2 2\phi + 0.000000024 \sin^2 \phi \sin^2 2\phi + \dots \\ &= 0.998321724 + 0.001676162 \cos 2\phi + 0.000002111 \cos 4\phi + 0.000000003 \cos 6\phi + \dots\end{aligned}\quad (199)$$

or

$$\begin{aligned}R &= 6378170.0 - 21489.7 \sin^2 \phi + 108.6 \sin^4 \phi - 0.6 \sin^6 \phi \pm \dots \\ &= 6378170.0 - 21381.7 \sin^2 \phi - 27.0 \sin^2 2\phi + 0.2 \sin^2 \phi \sin^2 2\phi + \dots \\ &= 6367465.7 + 10690.8 \cos 2\varphi + 13.5 \cos 4\varphi + 0.02 \cos 6\varphi + \dots\end{aligned}\quad (200)$$

and

$$\begin{aligned}\frac{R}{R_e} &= 1 - 0.003324310 \sin^2 \varphi - 0.000027777 \sin^4 \varphi - 0.000000241 \sin^6 \varphi \pm \dots \\ &= 1 - 0.003352330 \sin^2 \varphi - 0.000007004 \sin^2 2\varphi + 0.000000060 \sin^2 \phi \sin^2 2\phi + \dots \\ &= 0.998320349 + 0.001676156 \cos 2\varphi + 0.000003487 \cos 4\varphi + 0.000000008 \cos 6\varphi + \dots\end{aligned}\quad (201)$$

or

$$\begin{aligned}R &= 6378170.0 - 21203.0 \sin^2 \varphi - 177.2 \sin^4 \varphi - 1.5 \sin^6 \varphi \pm \dots \\ &= 6378170.0 - 21381.7 \sin^2 \varphi - 44.7 \sin^2 2\varphi + 0.4 \sin^2 \varphi \sin^2 2\varphi + \dots \\ &= 6367456.9 + 10690.8 \cos 2\varphi + 22.2 \cos 4\varphi + 0.05 \cos 6\varphi + \dots\end{aligned}\quad (202)$$

and

$$\begin{aligned}\frac{\mu}{\mu_e} = \frac{\rho_m}{\rho_{m_e}} &= 1 + 0.010040132 \sin^2 \varphi + 0.000084004 \sin^4 \varphi \\ &= 1.005051568 - 0.005062068 \cos 2 \varphi + 0.000010500 \cos 4 \varphi \mp \dots\end{aligned}\quad (203)$$

or

$$\begin{aligned}\mu &= 110574.95 + 1110.19 \sin^2 \varphi + 9.29 \sin^4 \varphi + \dots \\ &= 111133.53 - 559.74 \cos 2 \varphi + 1.16 \cos 4 \varphi \mp \dots\end{aligned}\quad (204)$$

$$\frac{\nu}{\nu_e} = \frac{\rho_n}{R_e} \cos \varphi = 1.000838785 \cos \varphi - 0.000839841 \cos 3 \varphi + 0.000001059 \cos 5 \varphi + \dots\quad (205)$$

or

$$\nu = 111413.44 \cos \varphi - 93.49 \cos 3 \varphi + 0.12 \cos 5 \varphi\quad (206)$$

and

$$\begin{aligned}\frac{\rho_n}{R_e} &= 1 + 0.003346711 \sin^2 \varphi + 0.000016801 \sin^4 \varphi + 0.000000094 \sin^6 \varphi + \dots \\ &= 1 + 0.003363605 \sin^2 \varphi - 0.000004224 \sin^2 2 \varphi - 0.000000023 \sin^2 \varphi \sin^2 2 \varphi \\ &= 1.001679685 - 0.001681800 \cos 2 \varphi + 0.000002118 \cos 4 \varphi - 0.000000003 \cos 6 \varphi\end{aligned}\quad (207)$$

or

$$\begin{aligned}\rho_n &= 6378170.0 + 21345.9 \sin^2 \varphi + 107.2 \sin^4 \varphi + 0.6 \sin^6 \varphi + \dots \\ &= 6378170.0 + 21453.6 \sin^2 \varphi - 26.9 \sin^2 2 \varphi - 0.15 \sin^2 \varphi \sin^2 2 \varphi + \dots \\ &= 6388883.3 - 10726.8 \cos 2 \varphi + 13.5 \cos 4 \varphi - \dots\end{aligned}\quad (208)$$

## REFERENCES

1. C. W. Allen: *Astrophysical Quantities*. 263pp., The Athlone Press, University of London, 1955.
2. G. W. Pettengill, H. W. Briscoe, J. V. Evans, E. Gehrels, G. M. Hyde, L. G. Kraft, R. Price, and W. B. Smith: *A Radar Investigation of Venus*. The Astronomical Journal, vol. 67, no. 4, pp. 181-190, May 1962.
3. D. O. Muhleman, D. B. Holdridge, and N. Block: *The Astronomical Unit Determined by Radar Reflections from Venus*. The Astronomical Journal, vol. 67, no. 4, pp. 191 - 203, May 1962.
4. K. D. Froome: *Precision Determination of the Velocity of Electromagnetic Waves*. Nature, vol. 181, no. 4604, p. 258, 25 Jan. 1958.
5. Simon Newcomb: *Tables of the Motion of the Earth on Its Axis and Around the Sun*. Astronomical Papers prepared for the use of the American Ephemeris and Nautical Almanac, vol. vi, part 1, 1895.
6. H. Spencer Jones: *The Rotation of the Earth, and the Secular Accelerations of the Sun, Moon and Planets*. Monthly Notices, Royal Astron. Soc., vol. 99, no. 7, pp. 541- 558, 1939.
7. G. M. Clemence: *On the System of Astronomical Constants*. The Astronomical Journal, vol. 53, no. 6, pp. 169-179, May 1948.
8. E. W. Brown: *Tables of the Motion of the Moon*. Yale University Press, New Haven, 1919.
9. Procès Verbaux des Séances du Comité International des Poids et Mesures, deuxième Série, tome 25, p. 77, 1957.
10. W. deSitter and Dirk Brouwer: *On the System of Astronomical Constants*. Bulletin of the Astronomical Institutes of the Netherlands, vol. VIII, no. 307, pp. 213-231, 8 July 1938.
11. Harold Jeffreys: *The Earth. Its Origins, History and Physical Constitution*. Cambridge University Press, Fourth Edition, 1959.
12. S. Herrick, R.M.L. Baker, Jr. and C. G. Hilton: *Gravitational and Related Constants for Accurate Space Navigation*. American Rocket Society Paper 497-57, October 1957.
13. M. W. Makemson, R.M.L. Baker, Jr. and G. B. Westrom: *Analysis and Standardization of Astrodynamical Constants*. Univ. of California, Los Angeles. Astrodynamical Report no. 12, February 1961. Journal of the Astronautical Sciences, vol. VIII, no. 1, Spring 1961; AAS Preprint 61-27, January 1961.
14. Simon Newcomb: *The Elements of the Four Inner Planets and the Fundamental Constants of Astronomy*. Washington, D.C., 1895 (Supplement to The American Ephemeris and Nautical Almanac for 1897).

REFERENCES (Continued)

15. Eugene Rabe: *Derivation of Fundamental Astronomical Constants from Observations of Eros During 1926-1945*. The Astronomical Journal, vol. 55, no. 1184, pp. 112-126, 1950.  
*Additional Note on the Solar Parallax from Eros*. The Astronomical Journal, vol. 59, pp. 409-411, 1954.
16. Julius Bauschinger: Enzykl. d. Math. Wiss. 6 II, no. 17, 1919. (The value is given in Ref. 45, p. 51).
17. H. Battermann: Veröffentl. d. Sternw. Koenigsberg/Pr. (The value is given in Ref. 45, p. 51)
18. Dirk Brouwer and O. B. Watts: Unpublished. (The value is given in Ref. 45, p. 52).
19. David Gill: *Determination of the Solar Parallax and the Mass of the Moon from Heliometer Observations of Iris, Victoria and Sappho Made in the Years 1888 and 1890*. Ann. Cape Observatory, vol. 6, 1896, and vol. 7, 1897.
20. A. R. Hinks: *Solar Parallax*. Monthly notices, Royal Astronomical Soc., vol. 70, no. 9, p. 588, 1909.
21. H. Spencer Jones: *The Solar Parallax and the Mass of the Moon from Observations of Eros at the Opposition of 1931*. Mem. Royal Astr. Soc., vol. 66, pt. 2, 56pp., London, 1941.
22. H. R. Morgan and F. P. Scott: *Observations of the Sun 1900-1937 Compared with Newcomb's Tables*. The Astronomical Journal, vol. 47, p. 193, 1939.
23. Harold Jeffreys: *On the Lunar Equation*. Monthly Notices of the Royal Astronomical Soc. (London), vol. 102, no. 4, pp. 194-204, 1942.
24. Erwin Delano: *The Lunar Equation from Observations of Eros 1930-31*. The Astronomical Journal, vol. 55, pp. 129-132, 1950.
25. Simon Newcomb: *Researches on the Motion of the Moon, Part II*. Astr. Paper. Amer. Eph., vol. IX, p. 39, 1912.
26. The American Ephemeris and Nautical Almanac for the Year 1962, U.S. Government Printing Office, Washington, D.C., 1960. Explanatory Supplement to the Astronomical Ephemeris and the American Ephemeris and Nautical Almanac, London, 1961.
27. H. Hirose and R. Manabe: *Motion and Figure of the Moon*. Transactions I.A.U., vol. 9, p. 265, 1955.
28. George H. Darwin: *The Theory of the Figure of the Earth, Carried to the Second Order of Small Quantities*. Monthly Notices Roy. Astron. Soc., vol. 60, pp. 82-124, 1899.
29. F. R. Helmert: *Die mathematischen und physikalischen Theorien der Höheren Geodäsie*. I. *Die mathematischen Theorien*. Leipzig, 1880. II. *Die physikalischen Theorien*. Leipzig, 1884.

REFERENCES (Continued)

30. E. C. Bullard: *The Figure of the Earth*. Monthly Notices Roy. Astron. Soc., Geophys. Suppl., vol. 5, no. 6, 1948.
31. W. deSitter: *On the Flattening and the Constitution of the Earth*. Bulletin of the Astronomical Institutes of the Netherlands, vol. II, no. 55, pp. 97-108, 14 May 1924.
32. F. R. Helmert: *Neue Formeln für den Verlauf der Schwerkraft im Meeresspiegel beim Festland*. Sitz-Ber. d. Kgl. Preuss. Akad. d. Wiss., vol. 41, pp. 676-68, Berlin 1915.
33. A. Berroth: *Die Erdgestalt und die Hauptträgheitsmomente A und B im Äquator aus Messungen der Schwerkraft*. Gerl. Beitr. Geophys., vol. 14, pp. 215-257, Leipzig, 1915-1918.
34. W. A. Heiskanen: *Untersuchungen über Schwerkraft und Isostasie*. Publ. Finn. Geod. Inst., no. 4, Helsinki, 1924.
35. W. A. Heiskanen: *Ist die Erde ein dreiachsiges Ellipsoid?* Gerlands Beitr. Geophys., vol. 19, pp. 356-377, Leipzig, 1928, Astr. Nachr., vol. 234, no. 5562, Kiel, 1928.
36. W. A. Heiskanen: *Investigations on the Gravity Formula*. Ann. Acad. Scient. Fennicae, series A, vol. 51, no. 8, Helsinki, 1938.
37. N. F. Shuravlev: *Die Bestimmung der Abplattung des Erdellipsoides aus gravimetrischen Messungen* (in Russian with English Abstract) Publ. Astro. Inst. Sternberg, vol. 14, pt. 2, Moscow, 1940.
38. Erkki Niskanen: *Gravity Formulas Derived by the Aid of the Level Land Stations*. Publ. Isos. Inst. IAG (Helsinki), no. 16, 1945; Ann. Acad. Scient. Fennicae, series A, no. III, 10, pp. 1-16, Helsinki, 1945.
39. Harold Jeffreys: *The Figures of the Earth and Moon* (Third Paper). Monthly Notices of the Roy. Astr. Soc., Geophys. Suppl., vol. 5, no. 7, pp. 219-247, July 1948.
40. H. Schütte: *Die Bestimmung der Abplattung der Erde aus der Schwerkraftverteilung nach dem Schwerereverzeichnis von N. F. Shuravlev*. Gerl. Beitr. Geophys., vol. 62, pp. 9-26, Leipzig, 1950.
41. U. A. Uotila: *Determination of the Shape of the Geoid*. Symposium: Size and Shape of the Earth, held at the Ohio State University, Columbus, Ohio, Nov. 13-15, 1956. Publ. no. 7, Inst. of Geodesy, Photogrammetry and Cartography, 1957.
42. W. A. Heiskanen: *Achievements of the World-wide Gravity Program of the Mapping and Charting Research Laboratory of the Ohio State University*. Report to the Toronto General Assembly of the IUGG in 1957.
43. F. Kühnen und Ph. Furtwängler: *Bestimmung der absoluten Größe der Schwerkraft zu Potsdam mit Reversionspendeln*. Veröff. d. Preuss. Geod. Inst. N.F. 27, 1906.
44. W. A. Heiskanen and F. A. Vening Meinesz: *The Earth and Its Gravity Field*. x + 470 pp., McGraw-Hill Book Co., New York, 1958 (p. 75).

REFERENCES (Continued)

45. Landolt-Börnstein: *Zahlenwerte und Funktionen aus Physik, Chemie, Astronomie, Geophysik und Technik*. Vol. III, pp. 43-52 (A. Kopff: Astrometrische Konstanten) and pp. 256-270 (K. Jun g: Schwerkraft und Erdfigur), Springer-Verlag, Berlin, 1952.
46. H. Andoyer: *Les formules de la précession après S. Newcomb*. Bulletin Astronomique (Paris), vol. 28, p. 67-76, 1911.
47. H. R. Morgan: *Systematic Corrections to the Albany General Catalogue*. The Astron. Jour., vol. 54, no. 1, pp. 1-9, Sept. 1948; *Corrections to the Fundamental Catalogs*. The Astron. Jour., vol. 54, no. 6, pp. 145-150, June 1949
48. Landolt-Börnstein: *Zahlenwerte und Funktionen*. (Ref. 45, p. 52.)
49. J. H. Oort: *The Constants of Precession of Galactic Rotation*. Bulletin of the Astron. Institutes of the Netherlands, vol. 9, p. 424 - 427, 1943.
50. R. R. Radau: *Sur la loi des densités à l'intérieur de la Terre*. Comptes Rendus Acad. Sci. (Paris), vol. 100, pp. 972-974, 1885; Bulletin Astronomique 2, p. 157, 1885.
51. O. Callandreau: *Sur le développement en série du potential*. J. Polytech., 1889; Ann. de l'Observ. de Paris 19 E, 1889.
52. Harold Jeffreys: *On the Figures of the Earth and Moon*. Monthly Notices of the Roy. Astr. Soc., vol. 97, no. 1, pp. 3-15, Nov. 1936 or Geophys. Suppl., vol. 4, no. 1, pp. 1-13, Jan. 1937.
53. Harold Jeffreys: *The Moon's Libration in Longitude*. Monthly Notices of the Roy. Astr. Soc., vol. 117, no. 5, pp. 475-477, 1957.
54. Harold Jeffreys: *On the Figure of the Moon*. Monthly Notices Roy. Astr. Soc., vol. 122, pp. 421-432, 1961.
55. W. M. Kaula: *A Geoid and World Geodetic System Based on a Combination of Gravimetric, Astrogeodetic, and Satellite Data*. Jour. Geophys. Res., vol. 66, no. 6, pp. 1799-1811, June 1961.
56. Victor C. Clarke Jr.: *Constants and Related Data Used in Trajectory Calculations at the Jet Propulsion Laboratory*. Jet Propulsion Laboratory, Technical Report No. 32-273, 14 pp., 1 May 1962.
57. Irene Fischer: *Parallax of the Moon in Terms of a World Geodetic System*. The Astronomical Jour., vol. 67, no. 6, pp. 373-378, Aug. 1962.
58. B. E. Kalensher: *Selenographic Coordinates*. Jet Propulsion Laboratory, Technical Report No. 32-41, 31 pp., 24 February 1961.

APPROVAL

ON A CONSISTENT SYSTEM OF ASTRODYNAMIC CONSTANTS

By

*Helmut G.L. Krause*

The information in this report has been reviewed for security classification. Review of any information concerning Department of Defense or Atomic Energy Commission programs has been made by the MSFC Security Classification Officer. This report, in its entirety, has been determined to be unclassified.

*Erich E. Goerner*

ERICH E. GOERNER  
Chief, Advanced Flight Systems Branch

*Wm A. Mrazek*

W. A. MRAZEK  
Director, Propulsion & Vehicle Engineering Division

## DISTRIBUTION

M-DIR	
Dr. von Braun	
M-DEP-R&D	
Dr. Rees	
M-CP-DIR	
Mr. Maus	
M-HME-P	
Dr. Rudolph	
M-AERO-DIR	
Dr. Geissler	
Dr. Hoelker	
M-AERO-TS	
Dr. Heybey	
Dr. Sperling	
M-AERO-PS	
Mr. Braunlich	
Mr. Vaughn	
Mr. Schmidt	
M-AERO-A	
Mr. Dahn	
M-AERO-D	
Mr. Horn	
Mr. Baker	
Mr. Hart	
Mr. Callaway	
Mr. Redus	
Mr. Thomae	
M-AERO-F	
Dr. Speer	
Mr. Kurtz	
M-AERO-G	
Mr. Dalton	
M-AERO-P	
Mr. Miner	
Mr. Braud	
Mr. Dearman	
Mr. Lisle	
Mr. Schmieder	
Mr. Schwaniger	
Mr. Silber	
Mr. Winch	
Mr. Davidson	
M-AERO-S	
Mr. de Fries	
M-ASTR-DIR	
Dr. Haeussermann	
	M-ASTR-TSS
	Mr. Kroh
	M-ASTR-A
	Mr. Digesu
	M-ASTR-I
	Dr. Decher
	M-ASTR-M
	Mr. Boehm
	Mr. Pfaff
	M-ASTR-F
	Mr. Hosenthien
	M-COMP-DIR
	Dr. Hoelzer
	Mr. Bradshaw
	M-COMP-TS
	Dr. Fehlberg
	Dr. Arenstorf
	M-FPO
	Mr. Koelle
	Mr. Williams
	Mr. Ruppe
	Mr. Gradecak
	M-L&M-DIR
	M-MS-H
	M-MS-I
	M-MS-IP
	M-MS-IPL (8)
	M-ME-DIR
	M-PAT
	M-P&VE-DIR
	Dr. Mrazek
	Mr. Weidner
	Mr. Hellebrand
	M-P&VE-V
	Mr. Palaoro
	M-P&VE-M
	Dr. Lucas
	M-P&VE-F
	Mr. Goerner
	Mr. Barker
	Mr. Swanson
	Dr. Krause (100)
	Mr. Burns (5)
	M-P&VE-FN
	Mr. Jordan
	Mr. Heyer
	M-HME-P

DISTRIBUTION (Continued)

M-P&VE-FN, Continued	M-SAT-DIR
Mr. Nixon	Dr. Lange
Mr. Saxton	M-TEST-DIR
Mr. Harris	Mr. Heimburg
Mr. Manning	
Mr. Whiton	Headquarters
Mr. Patton	National Aeronautics and Space Administration
M-P&VE-FF	Washington 25, D.C.
Mr. Galzerano	Dr. J. Shea, Dep. Dir.,
Mr. Fellenz	Manned Space Flight Office
Mr. Russell	
Mr. Akridge	Office of Scientific and Technical Information
Mr. Perry	National Aeronautics and Space Administration
Mr. Stafford	Washington 25, D.C.
Mr. Thomson	Acquisitions & Dissemination Branch (AFSS-AD)
Mr. Cohen	
Mr. Kromis	Ames Research Center
Mr. Croft	National Aeronautics and Space Administration
Mr. Messer	Moffett Field, California
Mr. Valentine	Library
Mr. Wheeler	Dr. Eggers
M-P&VE-FS	Goddard Space Flight Center
Mr. Pedigo	National Aeronautics and Space Administration
Mr. Neighbors	Greenbelt, Maryland
Mr. Laue	Library
Mr. Schwartz	Dr. J. A. O'Keefe
Mr. Orillion	Dr. J. W. Siry
Mr. Johns	Dr. W. M. Kaula
M-P&VE-P	Dr. W. D. Kahn
Mr. Paul	Dr. H. G. Hertz
Mr. Swalley	Dr. R. W. Bryant
M-P&VE-S	Dr. P. Musen
Mr. Kroll	
Dr. Glaser	Langley Research Center
M-P&VE-E	National Aeronautics and Space Administration
Mr. Schulz	Langley Field, Va.
M-QUAL-DIR	Library
Mr. Grau	Dr. J. C. Houbolt
M-RP-DIR	
Dr. Stuhlinger	Launch Operations Center
Mr. Heller	National Aeronautics and Space Administration
M-RP-N	Cape Canaveral, Florida
Dr. Shelton	Dr. K. Debus (2)
M-RP	Dr. A. Knothe
Dr. Schocken	Dr. Bruns
Mr. Naumann	
Mr. Snoddy	
Mr. Prescott	

DISTRIBUTION (Continued)

Lewis Research Center National Aeronautics and Space Administration 2100 Brookpark Road Cleveland 35, Ohio Library Dr. S. Himmel Dr. W. E. Moeckel	University of California, Dept. of Astronomy Los Angeles, California Library Dr. S. Herrick
Manned Spacecraft Center National Aeronautics and Space Administration Houston, Texas Library Mr. Faget	University of Cincinnati Observatory Cincinnati, Ohio Library Dr. P. Herget Dr. E. K. Rabe
Jet Propulsion Laboratory California Institute of Technology 4800 Oak Grove Drive Pasadena, California Library Mr. V. Clark	Lockheed Aircraft Corp. Burbank, California Library Dr. R. M. L. Baker
Air Force Cambridge Research Center L. G. Hanscom Field Bedford, Massachusetts Library Mr. O. W. Williams	Lockheed Aircraft Corp. Missile and Space Division Sunnyvale, California Library Dr. J. V. Breakwell Dr. A. F. Michielson Mr. W. B. Schramm
U. S. Naval Observatory Washington 25, D.C. Library Dr. G. M. Clemence, Dir.	North American Aviation Space and Information Systems Division Downey, California Library Dr. Laidlow
Yale Observatory Box 2023, Yale Station New Haven, Connecticut Library Dr. Dirk Brouwer, Dir.	Martin Company Space Systems Division, J-325 Baltimore 3, Maryland Library Dr. George E. Townsend (3)
Smithsonian Astrophysical Observatory Cambridge 38, Massachusetts Library Dr. Fred L. Whipple, Dir. Dr. Y. Kozai Dr. I. G. Izsak Dr. George Veis	General Dynamics/Astronautics San Diego, California Library Mr. F. Dore Dr. Krafft A. Ehricke
	Space Technology Laboratory, Inc. Redondo Beach, California Library Dr. P. Dergarabedian Dr. A. B. Michelwait

DISTRIBUTION (Continued)

Boeing Scientific Research Laboratories  
Seattle 24, Washington  
Library  
Dr. Angelo Miele

Chance Vought Corp.  
Astronautics Division  
Dallas 22, Texas  
Library  
Mr. Clarke, President

Douglas Aircraft Co.  
Missile and Space Systems Division  
3000 Ocean Park Boulevard  
Santa Monica, California  
Library

General Electric Co.  
Space Sciences Lab.  
3750 D Street  
Philadelphia 24, Pa.  
Library  
Dr. V. G. Szebehely

Aerospace Corp.  
El Segundo, California  
Library  
Dr. E. Levin

Republic Aviation Corp.  
Farmingdale, L.I., New York  
Library  
Mr. Samuel Pines  
Dr. Mary Payne

Rand Corp.  
Santa Monica, California  
Library  
Dr. Buchheim

Ballistic Research Laboratories  
Aberdeen Proving Ground, Maryland  
Library  
Dr. Boris Garfinkel

National Bureau of Standards  
Washington, D.C.  
Library  
Dr. John P. Vinti

Armed Services Tech. Information Agency (ASTIA)  
Arlington Hall Station  
Arlington 12, Virginia  
Library

Scientific Information Center  
Army Missile Command  
Redstone Arsenal, Huntsville, Alabama  
Library (5)

Scientific and Technical Information Facility (2)  
Attn: NASA Representative (S-AK/RKT)  
P. O. Box 5700  
Bethesda, Maryland

